## **CAMBRIDGE INTERNATIONAL EXAMINATIONS**

**International General Certificate of Secondary Education** 

## MARK SCHEME for the May/June 2014 series

## 0610 BIOLOGY

0610/51

Paper 5 (Practical Test), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2014 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



Page 2	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0610	51

Question	Mark Scheme		Marks	Comments		
1 (a)	lodine sol	ution or reagent/	iodine in KI;		I iodine alone	
	brown/or	ange/yellow to bl	ue/purple/black/AW;	[2]		
(b)	no reducing sugar present – remains blue;			A no change		
	low concentration – green/yellow;				A intermediate shades	
	high cond	entration – orang	e/red;	[3]	A brick red	
(c)	time / mins	starch test	reducing sugar		Check Supervisors Report.  both observations in each row ;;;	
	0	obs – concl	obs – concl. –		Both conclusions in each row match observations ;;;	
	5	obs – concl	obs – concl. –			
	10	obs – concl. –	obs – concl. –			
				[6]		
(d)	break down of starch; to form reducing sugar;			Check Supervisors Report.		
	reference to the candidate's own results;		max [3]			
(e) (i)	avoid contamination/prevent mixing/AW;		[1]			
(ii)	to see colour change easily or clearly/AW;		[1]			

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0610	51

(	(f)	replace enzyme with water; inactive enzyme by use of a low temperature/denature with high temperature;	[2]	
(	(g)	replication qualified/identify anomalies/AW;	[1]	
(	(h)	amylase/carbohydrase/maltase;	[1]	
(	(i) (i)	A – axis labelled and scaled <u>evenly;</u>		Accurate to $\pm$ 0.5 of small square.
		<ul> <li>S – size – plots for 'time' must use half or more of the axis;</li> <li>P – all points plotted accurately;</li> </ul>		
		L – line through all points;	[4]	
	(ii)	pH 7(.0);	[1]	
	(iii)	below optimum/pH 7or neutral as pH increases the activity increases/time decreases/AW; above optimum/pH 7 or neutral as pH increases the activity decreases/time increases/AW; credit use of figures;		To gain credit a comparison between two data points with a calculation should be shown.
		decreased activity/increased time occurs more rapidly/has steeper curve above pH 7;	max [3]	
			[Total: 28]	
2 (	(a)	(line and ) label/(i) to xylem of gorse;		
		(line and ) label/(ii) to phloem of gorse;	[2]	

Page 4	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2014	0610	51

(b)	in/from/via xylem (of gorse);	[1]	
(c)	measurement of <b>MN</b> : $9 \pm 1$ [mm];		A ecf for calculation
	formula : length ÷ 50;		
	actual width : 0.18 [mm]	[3]	
d (i)	O – outline – clear unbroken line and no shading;		
	<ul><li>S – size;</li><li>D – detail;</li><li>L – one correct label from: leg/limb/cephalohorax/</li></ul>		Drawing larger than 70 mm at widest point between legs.  A evidence of jointed leg(s) and mouth parts
	mouthpart;	[4]	
(ii)	Arachnid(a);		
	4 pairs or 8 legs/2 parts to body;	[2]	
		[Total: 12]	