## **CAMBRIDGE INTERNATIONAL EXAMINATIONS**

**International General Certificate of Secondary Education** 

## MARK SCHEME for the October/November 2013 series

## 0620 CHEMISTRY

0620/32

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October / November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



Page 2	ICCSE	Mark Scheme - October/November 201	Syllabus 13 0620	Paper 32
(a) Can	1	- October/November 20	13 0020	[1]
(b) A				[1]
(13)				(.)
(c) B				[1]
(d) D				[1]
(e) E				[1]
(f) A an	<b>d</b> D			[1]
				[Total: 6]
? (a) (i)	two atoms per moled	<u>cule</u>		[1]
(ii)	7e in outer shell or le	evel / same number of out	er electrons / need to gain or	ne electron [1]
(iii)	different number of e	energy levels / different nu	ımber of electrons	[1]
(iv)				_
	halogen	solid, liquid or gas at room temperature	colour	
	chlorine	gas	yellow / yellow green / green	
	bromine	liquid	<u>brown</u> / red- <u>brown</u> / orange- <u>brown</u> not: red / orange	
	iodine	solid	black / grey / silver-grey / purple / violet <b>NOT</b> : blue-black	
	NOTE: one mark for	each vertical column	L	[2]
	ect formula, AsF <sub>3</sub>	100		[1]
	os and 1bp around a and 1nbp around a			[1] [1]
		es / causes forward reaction	on / light causes	[4]
(incr		es the amount of silver (ar s reaction / uses up silver	nd so darkens glass) / silver reacts (and so reduc	[1] [1] es darkness)[1]
				[Total: 11]

Page 3	Mark Scheme	Syllabus	Paper	
	IGCSE – October/November 2013	0620	32	

3 (a) (i) the (forward) reaction is endothermic [1]

(ii) none volume of reactants and products the same **ACCEPT**: number of moles or molecules

[1] [1]

(iii) the reaction (between oxygen and nitric oxide) is exothermic [1] high temperatures push equilibrium to left / high temperatures decrease yield of products / low temperatures favour forward reaction [1]

(iv)  $4NO_2 + O_2 + 2H_2O \rightarrow 4HNO_3$ not balanced = (1) only

[2]

(v) (cost of) high amount of electricity / energy

[1]

(b) (i) contains more nitrogen

[1] [1]

(ii) photosynthesis chlorophyll is catalyst / chlorophyll absorbs light carbon dioxide and water react

[1] [1]

to make glucose / carbohydrates / starch / sugar / named sugar

[1]

[Total: 13]

(a) Any one of:

 $Fe_2O_3 + 3C \rightarrow 2Fe + 3CO$  $2Fe_2O_3 + 3C \rightarrow 4Fe + 3CO_2$  $Fe_2O_3 + 3CO \rightarrow 2Fe + 3CO_2$ for correct equation (2) not balanced = (1) only

any four of:

coke burns to form carbon dioxide / C +  $O_2 \rightarrow CO_2$ 

this reacts with more carbon to form carbon monoxide / C +  $CO_2 \rightarrow 2CO$ 

calcium carbonate decomposes to form calcium oxide and carbon dioxide / CaCO₃ → CaO +  $CO_2$ 

calcium oxide / calcium carbonate reacts with silica / silicon oxide / silicon(IV) oxide (in ore) to form calcium silicate / slag / CaO + SiO<sub>2</sub>  $\rightarrow$  CaSiO<sub>3</sub> or CaCO<sub>3</sub> + SiO<sub>2</sub>  $\rightarrow$  CaSiO<sub>3</sub> + CO<sub>2</sub>

the reaction between carbon and oxygen is exothermic / produces heat / coke is used as a fuel / the slag floats on the (molten) iron / the slag and molten iron can be run off separately

[6]

		IGCSE – October/November 2013	0620	32
(b) (i	, .	enhouse effect / CO <sub>2</sub> is a greenhouse gas al warming / ice caps melting / suitable example		[1] [1]
(ii	(ii) burning or combustion of charcoal produces carbon dioxide trees use carbon dioxide (in photosynthesis)		[1] [1]	
(iii	) cath	ode reaction Fe³+ 3e → Fe		[1]
		de reaction 2O²- → O₂ + 4e palanced = (1) only		[2]
				[Total: 13]
` ,	ecause harges	they have more than one oxidation state or va	alency / form ior	ns with different [1]
		e two iron oxides (iron(III) oxide and iron(II) ox nds / iron forms iron(II) and iron(III) compounds	ide) / iron forms	Fe <sup>2+</sup> and Fe <sup>3+</sup> [1]
(b) (i	) to re	emove the precipitate / remove the silver(I) chromate	e(VI) / remove the	residue [1]
(ii	•	emove <u>soluble</u> impurities / remove named <u>soluble</u> sa tants	alt e.g. potassium	nitrate / remove [1]
(iii	) to dr	y solid / to remove water		[1]
(c) (i	•	d <u>one</u> mole of potassium chromate(VI) for <u>two</u> nences to mole ratio	noles of silver(I)	nitrate / correct [1]
(ii	NO1	s of AgNO <sub>3</sub> needed is 170 × 0.2 × 0.1 = 3.4g E: if answer given is 34 they have omitted 0.1 <b>OW</b> : (1) ecf		[2]
(iii	) num	ber of moles of AgNO <sub>3</sub> used = $0.02 \times 0.2 = 0.004$		[1]
	num	ber of moles of Ag <sub>2</sub> CrO <sub>4</sub> formed = 0.002		[1]
	mas	s of one mole of $Ag_2CrO_4 = 332g$		
		s of Ag₂CrO₄ formed = 0.664g E: use ecf when appropriate		[1]
				[Total: 11]

Mark Scheme

Syllabus

Paper

[Total: 11]

Page 4

		<b>3</b>	IGCSE – October/November 2013 0620 32	
6	(a)	(i)	$Cu(OH)_2 \rightarrow CuO + H_2O$	[1]
		(ii)	Rb	[1]
	(b)	(i)	electron loss	[1]
		(ii)	because they can accept electrons	[1]
	(c)	(i)	copper and mercury	[1]
		(ii)	add copper / mercury / metal to (named) acid <b>and</b> no reaction / no bubbles / hydrogen	no [1]
	(d)	(i)	Mn	[1]
		(ii)	(solution) becomes colourless / decolourises NOT: clear	[1]
			[Total	: 8]
7	(a)	(i)	contains <u>only</u> carbon, hydrogen and oxygen hydrogen (atom) to oxygen (atom) ratio is 2:1 <b>ALLOW</b> : C:H:O as 1:2:1 or C <sub>n</sub> (H <sub>2</sub> O) <sub>n</sub>	[1] [1]
		(ii)	condensation polymerisation	[1] [1]
	(b)	(i)	cells / micro-organisms / plants / animals / metabolic reactions obtaining energy from food / glucose / nutrients	[1] [1]
		(ii)	$2C_2H_5OH + 2CO_2$ allow: $C_2H_6O$ for $C_2H_5OH$ not balanced = (1) only	[2]
		(iii)	to prevent aerobic respiration / to get anaerobic respiration / to prevent ethanoic aclactic acid / carboxylic acids being formed / to prevent oxidation of ethanol	id / [1]
	(c)	NO	ayed formula of methyl butanoate E: all bonds must be shown E: award (1) if error in alkyl groups but correct displayed structure of –COO–	[2]
	(d)	(i)	alcohol, e.g. glycerol, circled <b>ALLOW</b> : if only part of glycerol molecule is circled as long as it involves an OH group	[1]
		(ii)	saturated correct reason based on group $C_{17}H_{35}$ / all C–C bonds / no C = C bonds	[1]

Mark Scheme

Syllabus

Paper

Page 5

Page 6	Mark Scheme	Syllabus	Paper
	IGCSE – October/November 2013	0620	32

1	(iii) salt / carboxylate / alkanoate (making) soap ACCEPT: detergent / washing	[1] [1]
(e)	at least one correct amide linkage –CONH– continuation shown at both ends of chain diagram showing three (different) amino acid residues	[1] [1] [1]

[Total: 18]