CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International General Certificate of Secondary Education

www.PapaCambridge.com MARK SCHEME for the October/November 2014 series

0620 CHEMISTRY

0620/21

Paper 2 (Core Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

BB CAMBRIDGE

	logo '	<u>. </u>	Mark Scheme	Si 2. D
	Page 2	_	Cambridge IGCSE – October/November 2014	062 W Oer
4	/ ₂ \	/:\		25
1	(a)	(i)	E	THE
		(ii)	A <u>and</u> D	Sylvar day per 062 Office of the sylvar of t
		(iii)	D	[1]
		(iv)	В	[1]
		(v)	D	[1]
		(vi)	A and D	[1]
	(b)	C ₂ ŀ	H_4Br_2	[1]
	(c)	4 (I	H ₂ O)	[1]
		5 (0	O ₂) te: mark dependent on 4 (H ₂ O)	[1]
				[Total: 9]
2	(a)	(i)	sodium / Na ⁺	[1]
		(ii)	X is fluoride	[1]
			Y is nitrate	[1]
		(iii)	0.244 (mg) allow : 0.24	[1]
		(iv)	4th box down ticked (weakly acidic)	[1]
	(b)	(ad	d nitric acid) add silver nitrate	[1]
			te precipitate te: mark dependent on correct reagent	[1]
	(c)	pol	ymer	[1]
		mo	nomer	[1]

[Total: 9]

		Cambridge IGCSE – October/November 2014 062		
(a)	ring	Cambridge IGCSE – October/November 2014 g around the OH group mine (water)	Morio	
(b)	bromine (water) allow: bromination		3	
	decolourised / turns colourless note: mark dependent on correct reagent ignore: goes clear / gets discoloured		[1]	
	allow : potassium manganate(VII) / potassium permanganate (1) turns colourless (1)			
	ign	ore: incorrect colour of reagent		
(c)	(i)	to break up the cells / to extract the pigment / to separate the pigment from the petals / idea of getting the colour out of the petals, e.g. otherwise the colour won't come out	[1]	
		idea that solvent dissolves the pigment / idea of making a solution ignore : find out how pure the rose petals are / reference to separating colours	[1]	
	(ii)	pigment might be absorbed onto filter paper / pigment sticks to filter paper	[1]	
(d)	(i)	chromatography	[1]	
	(ii)	spot near the bottom and above the solvent level	[1]	
((iii)	to keep atmosphere in jar saturated (with solvent vapour) allow: to reduce / prevent (solvent) evaporation	[1]	
((iv)	A and C	[1]	
(e)	(e) structure of ethanol with ALL atoms and bonds shown [2]			

Mark Scheme

Page 3

3

[Total: 12]

per

age 4	wark scheme	Syl
	Cambridge IGCSE – October/November 2014	062
(a)	thermometer	062 PARCAMBATOR
(b)	Any two from:	36
	 same volume of water in can same height of burner (from can) wick same height same rate / amount of stirring of water allow: same temperature of water at start allow: same amount of fuels burnt / same temperature rise allow: same type of can 	
(c)	so same temperature throughout the water / to stop differences in temperature parts of the water / otherwise the temperature will be higher bottom (of the water) / so not hotter in one place ignore: to mix the water / so there are no convection currents	
(d)	decreases / goes down	[1]
	idea of liquid or fuel turning to vapour / gas; allow: gases formed ignore: fuels evaporate note: 2nd mark dependent on first	[1]
(e)	F	[1]
(f)	(i) mixture of metals / mixture of metal(s) + non-metals do not allow: compound	[1]
	(ii) covers surface / idea of protective layer	[1]
	prevents contact with air / prevents contact with water / so air (or w no react with steel do not allow: reference to tin being more reactive / sacrificial protesecond marking point)	[1]
(g)	1st box down ticked (giant covalent)	[1]

Mark Scheme

Page 4

[Total: 11]

	<u> </u>		Cambridge IGCSE – October/November 2014 062	Day	
5	(a)	Any	y four from:	Sandy.	
		•	suitable named metal / metal oxide e.g. reactive metal such as Mg / Zn or their oxides suitable named acid metal + acid gives metal salt / named metal gives named metal salt metal + acid gives off hydrogen te: complete word equation for metal + acid → salt + hydrogen (2) metal oxide + acid gives metal salt / named metal oxide gives named metal salt water also product of reaction of metal oxide + acid te: complete word equation for metal oxide + acid → salt + water (2)	De Cambridge C	
	(b)	exc	othermic	[1]	
	(c)	thic	suitable use of radioactive isotope e.g. detecting leaks in pipes / checking thickness of paper / tracer / cancer treatment / investigating thyroid function ignore : atomic bombs / explosions		
	(d)	pro	tons 92 and 92	[1]	
		neu	utrons 143 and 146	[1]	
		ele	ctrons 92 and 92	[1]	
				[Total: 9]	
6	(a)	(i)	(concentration) decreases	[1]	
			then remains constant allow: levels out	[1]	
		(ii)	3.8 (hr) / 3 hr 48 min	[1]	
		(iii)	9 (hr) allow : 8.8–9.2 (hr)	[1]	
		(iv)	steeper graph line from same starting point	[1]	
			levels off lower than 0.10 mol /dm ³	[1]	

(v) increase the temperature / increase concentration of sodium hydroxide allow: add a catalyst

Mark Scheme

Page 5

5

6

[1]

per

Mark Scheme Syl	per
Cambridge IGCSE – October/November 2014 062	No.
Any four from:	Camp
 acid in burette use (volumetric) pipette to put sodium hydroxide into flask allow: sodium hydroxide in burette / acid in flask idea of correct setup of apparatus, i.e.flask under burette indicator in flask run hydrochloric acid into sodium hydroxide until indicator changes colour any indication of good technique e.g. repeating experiment / add acid slowly / shaking flask after each addition of acid note: answers must be in the correct context, e.g. do not allow indicator in burette 	Tag
bonding pair of electrons between H and Cl and no additional electrons on the H atom six non-bonding electrons around the chlorine atom ignore : inner shell electrons in Cl .	[1] [1]
	[Total: 13]
or minerals) lost from soil when crops harvested / for more plant protein allow : to give more nutrients to plants	[1]
neutralisation acid-base (reaction)	[1]
ammonium nitrate	[1]
2 NH_4^+ to 1 SO_4^{2-} / 2 ammonium to 1 sulfate allow : 2:1 or 1:2 ratio unqualified allow : $(NH_4)_2SO_4$	[1]
 Any two from: slaked lime can form an alkaline solution with water / slaked lime is calcium hydroxide / slaked lime is a hydroxide / slaked lime is basic slaked lime reacts with ammonium (salts) 	[2]
	Any four from: acid in burette use (volumetric) pipette to put sodium hydroxide into flask allow: sodium hydroxide in burette / acid in flask idea of correct setup of apparatus, i.e.flask under burette indicator in flask run hydrochloric acid into sodium hydroxide until indicator changes colour any indication of good technique e.g. repeating experiment / add acid slowly / shaking flask after each addition of acid note: answers must be in the correct context, e.g. do not allow indicator in burette bonding pair of electrons between H and Cl and no additional electrons on the H atom six non-bonding electrons around the chlorine atom ignore: inner shell electrons in Cl. for better crop / for better plant growth / to replace elements (or named elements or minerals) lost from soil when crops harvested / for more plant protein allow: to give more nutrients to plants ignore: for healthy plant growth / to give plants the compounds they need to grow / to help plants grow neutralisation acid-base (reaction) ammonium nitrate 2 NH ₄ ⁺ to 1 SO ₄ ²⁻ / 2 ammonium to 1 sulfate allow: 2:1 or 1:2 ratio unqualified allow: (NH ₄) ₂ SO ₄ Any two from: slaked lime can form an alkaline solution with water / slaked lime is calcium hydroxide / slaked lime is a hydroxide / slaked lime is basic

ammonia escapes from soil / gas escapes from soil

Page	2 7	Mark Scheme Syl	per
90	•	Cambridge IGCSE – October/November 2014 062	da l
(f)) po	Mark Scheme Cambridge IGCSE – October/November 2014 sitive: anode and negative cathode + electrode → chlorine	Cally
	at	+ electrode → chlorine	Tage
	at	– electrode → potassium	[1]
			[Total: 9]
8 (a	ı) Ar	ny four from:	[4]
	•	dissolving diffusion in iodine solid the particles are close together in iodine solid the particles <u>only</u> vibrate ALLOW: particles do not move in solution the iodine molecules are further / far apart in solution the particles are randomly arranged/ no particular arrangement in solution, particles move (fairly) freely / in solution particles slide over solvent molecules ow: in solution particles move slowly (from place to place) in solution there is bulk movement of particles from higher to lower concentration / particles spread out in solution / move everywhere / mix up ow: particles move from higher to lower concentration ideas of explanation of dissolving in terms of solvent molecules getting between the iodine particles ideas about forces between particles of iodine being weakened on dissolving	
(b) (i)	solid	[1]
	(ii)	heat causes astatine to melt / energy causes astatine to melt allow:: the astatine has melted / radioactivity melts the astatine	[1]
	(iii)	At ₂ on right	[1]
		2 (NaAt) on left note : 2nd mark dependent on At ₂ or 2At on right	[1]

[Total: 8]