

---

**CHEMISTRY****0620/42**

Paper 4 Extended Theory

**October/November 2017**

MARK SCHEME

Maximum Mark: 80

---

**Published**

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2017 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

© IGCSE is a registered trademark.

This syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

---

This document consists of **7** printed pages.

Question	Answer	Marks
1(a)(i)	Brownian (motion)	1
1(a)(ii)	molecules	1
	nitrogen / N <sub>2</sub> / N <b>OR</b> oxygen / O <sub>2</sub> / O	1
1(a)(iii)	nitrogen <b>OR</b> oxygen (particles) collide with / bombard / hit the dust (particles)	1
	(the bombarding particles) move <b>randomly</b>	1
1(b)(i)	diffusion	1
1(b)(ii)	Br <sub>2</sub> has an <i>M<sub>r</sub></i> of 160 <b>AND</b> Cl <sub>2</sub> has an <i>M<sub>r</sub></i> of 71 / bromine has an <i>A<sub>r</sub></i> of 80 <b>AND</b> chlorine has an <i>A<sub>r</sub></i> of 35.5	1
	(heavier) bromine (molecules / particles) diffuses more slowly	1
1(b)(iii)	particles have more energy / move faster	1

Question	Answer	Marks
2(a)	Si: 2: 8 : 4	1
	Ca <sup>2+</sup> : 2 : 8: 8	1
	N <sup>3-</sup> : 2 : 8	1
2(b)	Ca <sub>3</sub> N <sub>2</sub>	1
2(c)	Li shown as having one shell with 2 electrons <b>OR</b> no electrons <b>OR</b> no outer shell	1
	Cl shown as having an outer shell of 7 electrons of one type, plus one different electron which matches Li electrons	1
	'+' charge on Li <b>AND</b> '-' charge on Cl	1

Question	Answer	Marks
2(d)	two shared pairs of electrons	1
	both <i>Cl</i> with complete outer shells	1
	S with complete outer shell	1
2(e)	$\text{SCl}_2$ has intermolecular forces (of attraction)	1
	$\text{LiCl}$ has (electrostatic) forces (of attraction) between ions	1
	intermolecular forces are weaker / less energy is needed to break intermolecular forces	1
2(f)	silicon(IV) oxide	1

Question	Answer	Marks
3(a)	<i>exothermic mark</i> : horizontal line representing the energy of the products below the energy of the reactants	1
	<i>label of products mark</i> : product line labelled with $2\text{CO}_2 + 3\text{H}_2\text{O}$	1
	<i>correct direction of vertical heat of reaction arrow</i> : arrow starts level with reactant energy and finishes level with product energy <b>AND</b> has (only) <b>one</b> arrow head	1
3(b)	activation energy / $E_a$	1

Question	Answer	Marks
3(c)	<p>–650 kJ / mol</p> <p><b>M1</b> bonds broken</p> $2 \cdot ((3 \cdot 410) + 360 + 460) + (3 \cdot 500)$ $2 \cdot (1230 + 360 + 460) + 1500$ $2 \cdot 2050 + 1500$ $4100 + 1500 = 5600$ <p><b>M2</b> bonds formed</p> $(2 \cdot (2 \cdot 805)) + (4 \cdot (2 \cdot 460))$ $2 \cdot 1610 + 4 \cdot 920$ $3220 + 3680 = 6900$ <p><b>M3 = M1 – M2</b></p> <p>energy change of reaction = 5600 – 6900 = –1300</p> <p><b>M4 = M3 / 2</b></p>	4
3(d)(i)	cracking	1
3(d)(ii)	$\text{C}_{12}\text{H}_{26} \rightarrow 3\text{C}_2\text{H}_4 + \text{C}_6\text{H}_{14}$ <p><b>M1</b> <math>\text{C}_{12}\text{H}_{26}</math></p> <p><b>M2</b> rest of equation</p>	2
3(d)(iii)	phosphoric acid	1
	heat	1
3(d)(iv)	addition / hydration	1
3(d)(v)	measure its boiling temperature	1
	compare to (known) data	1
3(e)(i)	<p>any 2 from:</p> <p>37 °C</p> <p>anaerobic</p> <p>glucose is aqueous</p> <p>yeast</p>	2

Question	Answer	Marks
3(e)(ii)	$C_6H_{12}O_6 \rightarrow 2C_2H_5OH + 2CO_2$ <b>M1</b> CO <sub>2</sub> as a product <b>M2</b> Rest of equation	2
3(e)(iii)	yeast is killed by the ethanol	1
3(e)(v)	slow rate of reaction	1
3(e)(v)	uses renewable resources / does not use a finite resource	1
3(e)(vi)	fractional distillation	1
3(f)(i)	CH <sub>3</sub> O	1
3(f)(ii)	no (C=C) double bonds	1
3(f)(iii)	at least two alternating rectangles with attempted linking	1
	<b>one</b> displayed ester link (all atoms and all bonds)	1
	fully correct structure with at least one repeat unit including continuation bonds from correct atom or rectangle	1
3(f)(iv)	polyester	1

Question	Answer	Marks
4(a)	the breakdown (into elements)	1
	of an (ionic) compound by (the passage of) electricity	1
4(b)(i)	oxygen	1
4(b)(ii)	<b>glowing</b> splint	1
	relights	1

Question	Answer	Marks
4(b)(iii)	$2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2$ <b>M1</b> gain of electrons by $\text{H}^+$ <b>M2</b> rest of equation	2
4(c)	<i>the wires:</i> electrons	1
	<i>the electrolyte:</i> ions	1
4(d)	any 2 from: green gas at positive electrode bulb is brighter rate of bubbles increases	2
4(e)(i)	<i>anode made of:</i> impure copper	1
	<i>cathode made of:</i> (pure) copper	1
	<i>electrolyte of:</i> (aqueous) copper sulfate	1
4(e)(ii)	silver (impurities) fall to the bottom of the cell	1
	zinc (impurities) (dissolve) into solution (as ions)	1
	because zinc is more reactive than copper <b>AND</b> silver is less reactive than copper	1

Question	Answer	Marks
5(a)	both colours referred to correctly as observations in both parts of the answer	1
	(if sulfuric acid is added to solution Y,) equilibrium moves to the right-hand side	1
	because the concentration of acid has increased	1
	(if sodium hydroxide is added to solution Y,) equilibrium moves to the left-hand side	1
	because sodium hydroxide reacts with / neutralises sulfuric acid	1
5(b)(i)	210 cm <sup>3</sup> <b>M1</b> expected volume of hydrogen = 300 cm <sup>3</sup> <b>M2</b> 70% of <b>M1</b>	2
5(b)(ii)	fewer moles / molecules / particles (of gas) on the left-hand side	1
5(b)(iii)	endothermic	1
5(b)(iv)	increases rate (of reaction)	1