



## **Cambridge International Examinations**

Cambridge International General Certificate of Secondary Education

CHEMISTRY			062	20/41
CENTRE NUMBER		CANDIDATE NUMBER		
CANDIDATE NAME				

Paper 4 Theory (Extended)

October/November 2018

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of 15 printed pages and 1 blank page.



1 The following formulae represent different substances.								
		Al	Ag	CaCO <sub>3</sub>	CH₄	$Cl_2$	Cu	SO <sub>2</sub>

Answer the following questions using only these substances. Each substance may be used once, more than once or not at all.

State which substance is:

(a)	used to make food containers	[1]
(b)	added to a blast furnace to remove impurities during the production of iron	[1]
(c)	the main constituent of natural gas	[1]
(d)	a cause of acid rain	[1]
(e)	a gas which bleaches damp litmus paper	[1]
(f)	a gas which contributes to climate change.	[1]
	[Total	: 6]


2 The table gives some information about four different particles, **A**, **B**, **C** and **D**.

particle	number of electrons	number of neutrons	number of protons	electronic structure	charge on particle
Α	11	12	11	2,8,1	0
В		14	11	2,8,1	0
С	18	20		2,8,8	0
D	18	20	17		

(a)	Complete the table. The first row has been done for you.	[4]
(b)	Give <b>two</b> particles from the table which are isotopes of each other.	
		[1]
(c)	Element <b>Z</b> is in the same group of the Periodic Table as <b>A</b> and is less reactive than <b>A</b> .	
	State the identity of element <b>Z</b> .	
		[1]
(d)	C is unreactive.	
	Use information from the table to explain why.	
		[1]
	[Total	: 7]

3 (a) Copper(II) nitrate decomposes when heated. Two gases, oxygen and nitrogen dioxide, and a solid are made in the reaction.

A sample of copper(II) nitrate was decomposed using the apparatus shown.



(i) Complete the chemical equation for the reaction.

$$2Cu(NO_3)_2 \rightarrow O_2 + .....NO_2 + .....$$
 [2]

(ii) Only oxygen gas is collected at X.

Explain why.	
	[1]

(b) Nitrogen dioxide and other oxides of nitrogen are formed in car engines.

Explain how nitrogen dioxide is formed in car engines.

			[2]

(c) Ate	eacher he	eated 18.8g of copper	r(II) nitrate.						
(i)	Calculate the number of moles of copper(II) nitrate present in the 18.8 g.								
<i></i>	<b>.</b>								
(ii)	Calculate the maximum number of moles of oxygen that can be made by heating 18.8 g of copper(II) nitrate.								
				mol [1]					
(iii)			ne of oxygen at room temperature and pressure g of $copper(\mathrm{II})$ nitrate.	, in cm³, that					
				cm³ [1]					
(d) Asa	ample of	copper(II) nitrate was	s dissolved in water to form an aqueous solution	1.					
	e aqueou shown.	s solution was split in	to three portions. A separate test was done on	each portion					
	test	reagent added	result						
	1	aqueous sodium hydroxide	light blue precipitate forms						
	2	zinc powder	solution changes from blue to colourless and a brown solid forms						
	3		ammonia gas is produced						
(i)	Give the	e formula of the light b	olue precipitate formed in test 1.						
				[1]					
(ii)	Explain	the changes seen in t	test 2.						
				[3]					
(iii)	Identify	the <b>two</b> reagents that	t must be added to the aqueous copper(II) nitra	te in test 3.					
	1								

[2]

(e)		Copper(II) nitrate can be made by reacting copper(II) carbonate with nitric acid. One of the products is carbon dioxide.					
	(i)	Write a chemical equation for the reaction of copper(II) carbonate with nitric acid.	101				
			[2]				
	(ii)	Carbon dioxide is added to the air by living things.					
		Name the chemical process by which living things add carbon dioxide to the air.					
			[1]				
			[,]				
(	(iii)	Carbon dioxide is removed from the air by plants.					
		Name the chemical process by which plants remove carbon dioxide from the air.					
			[1]				
		[Total: 1	19]				

4	(a) Sul	furic acid is made industrially by a four-step process.	
	ste ste	<ul> <li>p 1 Sulfur is burned in air to produce sulfur dioxide.</li> <li>p 2 Sulfur dioxide is converted into sulfur trioxide.</li> <li>p 3 Sulfur trioxide is reacted with concentrated sulfuric acid to produce oleum.</li> <li>p 4 Oleum is reacted with water to produce concentrated sulfuric acid.</li> </ul>	
	(i)	Some sulfur is obtained by mining.  Name <b>one</b> other major source of sulfur.	
			[1]
	(ii)	What is the name of the process by which sulfuric acid is made industrially?	
	(iii)	Describe the conversion of sulfur dioxide into sulfur trioxide in <b>step 2</b> .	[1]
		In your answer, include:     a chemical equation for the reaction     the essential reaction conditions.	
			[5]
		en concentrated sulfuric acid is added to glucose, $C_6H_{12}O_6$ , a black solid is producencentrated sulfuric acid acts as a dehydrating agent.	ed. The
	(i)	What is removed from the glucose in this reaction?	
			[1]
	(ii)	Name the black solid produced in this reaction.	
			[1]

(c)		The gas hydrogen sulfide, $\rm H_2S$ , is produced when concentrated sulfuric acid is added to solid potassium iodide.				
	The	he reaction involves oxidation.				
	(i)	Define the term oxidation in terms of electron transfer.				
		[	[1]			
	(ii)	Complete the dot-and-cross diagram to show the electron arrangement in a molecule hydrogen sulfide. Show outer shell electrons only.	of			
		H S H	2]			
	(iii)	Hydrogen sulfide has a simple molecular structure.				
		Explain why hydrogen sulfide has a low boiling point.				
		[	2]			

1	d)	Dilute sulfuric acid	reacts with ac	meous sodium	hydrogencari	honate in a n	eutralisation i	reaction
1	u,	Dilute Sulluite acid	Teacis with at	lacons somini	rryurogencan	bonate in a n	Guliansanoni	Cacilon.

$$H_2SO_4(aq) + 2NaHCO_3(aq) \rightarrow Na_2SO_4(aq) + 2H_2O(I) + 2CO_2(g)$$

In a titration, 0.200 mol/dm³ aqueous sodium hydrogencarbonate was used to neutralise 20.0 cm³ of dilute sulfuric acid of concentration 0.150 mol/dm³.

(i)	Calculate	the nu	ımber d	of moles	of dilute	e sulfuric	acid	used in	the	titration.
-----	-----------	--------	---------	----------	-----------	------------	------	---------	-----	------------

			. mol	[1]

- (ii) Calculate the number of moles of sodium hydrogencarbonate needed to neutralise the dilute sulfuric acid.
  - ..... mol [1]
- (iii) Calculate the volume, in cm³, of 0.200 mol/dm³ aqueous sodium hydrogencarbonate needed to neutralise the dilute sulfuric acid.

..... cm<sup>3</sup> [1]

[Total: 17]

**5** Hydrogen gas reacts with iodine gas. The equation is shown.


$$H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$$

The reaction is reversible and can reach equilibrium.

(a) What is meant by the term equilibrium?

 	 ro.

**(b)** The graphs show how pressure affects the yield of hydrogen iodide, HI, at two different temperatures.




(i)	Explain why	the yield at	500°C does	<b>not</b> change as	s the pressu	re is increased
-----	-------------	--------------	------------	----------------------	--------------	-----------------

		[1]

(ii) What can you conclude from the difference in the yield of hydrogen iodide at the **two** temperatures shown? Explain your answer.

temperatures energy. Explain your anewer.	
	[2]

(c) The graph shows how the concentration of hydrogen iodide, HI, changes after hydrogen gas and iodine gas are mixed together in a sealed container.



(i)	When is the rate of reaction fastest?	
		[1

(ii) The reaction was repeated at the same temperature and pressure but in the presence of a catalyst.

Draw a graph on the same axes to show how the concentration of hydrogen iodide changes with time in the presence of a catalyst. [2]

(d) A mixture of hydrogen gas and iodine gas is allowed to reach equilibrium.

(i) Increasing the pressure of a gas increases its concentration.

State and explain the effect of increasing the pressure on the **rate** of the forward reaction.

(ii) State and explain the effect of increasing the temperature on the **rate** of the reverse reaction.

[Total: 13]

(a)		ane, $\rm C_2H_6$ , is a member of the homologous series called alkanes. anol, $\rm C_2H_5OH$ , is a member of the homologous series called alcohols.	
	(i)	Alkanes are hydrocarbons.	
		What is meant by the term hydrocarbon?	
		[	[2]
	(ii)	All members of a homologous series can be represented by a general formula.	
		State the general formula of:	
		alkanes	
		• alcohols	
	····\		[2]
(	(iii)	State <b>two</b> characteristics, other than having the same general formula, of members of homologous series.	а
		1	
		2	
		г	
		ι	[2]
(b)	Eth	ane can react with chlorine in a substitution reaction.	
	(i)	State <b>one</b> essential reaction condition.	
		[	[1]
	(ii)	Draw the structure of the organic product formed by substitution of <b>one</b> of the hydroge atoms in ethane with chlorine. Show all of the atoms and all of the bonds.	∍n
		[	[1]
(	(iii)	Name the product of the substitution reaction between ethane and chlorine that does no contain carbon.	ot
		[	[1]

6

(c) Propan-1-ol is an alcohol.

The structure of propan-1-ol is shown.

Propan-1-ol reacts with ethanoic acid to form an ester.

Give the name of the ester formed in this reaction.

......[1]

(d) Ester Y has the structure shown.

ester Y

(i)	Give	the	molecular	formula	of	ester	Y.	
-----	------	-----	-----------	---------	----	-------	----	--

.....[1]

(ii) Draw the structures of the carboxylic acid and the alcohol used to make ester Y. Show all of the atoms and all of the bonds. Give the name of the carboxylic acid and the alcohol.

structure of the carboxylic acid

name of the carboxylic acid .....

structure of the alcohol

name of the alcohol .....

[4]

(e)	Nylon is a polyamic	de.			
	Complete the diag bonds present in the	ram to show the structue linkages.	ure of nylon. Show al	I of the atoms and all	of the
					[3]

[Total: 18]

## **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The Periodic Table of Elements

₹	2 H	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton 84	25	Xe	xenon 131	98	R	radon			
₹			6	ш	fluorine 19	17	Cl	chlorine 35.5	35	B	bromine 80	53	н	iodine 127	85	¥	astatine -			
5																		116		ermorium -
>	_																			≧
≥								<u> </u>										114	Εl	erovium -
=																				
								<u>a</u>										112	C	pernicium
																				darmstadtium roe
5																				_
		uago																		
		hydro 1																		
						1			25	Mn	mangane 55	43								
			_	pol	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
		Key	atomic numbe	mic sym	name ative atomic m				23	>	vanadium 51	41	g	niobium 93	73	٦	tantalum 181	105	Op	dubnium —
				ato	rela				22	ı=	titanium 48	40	Zr	zirconium 91	72	士	hafnium 178	104	꿆	rutherfordium -
						1			21	Sc	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89–103	actinoids	
=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ഗ്	strontium 88	56	Ba	barium 137	88	Ra	radium -
_			3	:=	lithium 7	1	Na	sodium 23	19	¥	potassium 39	37	Rb	rubidium 85	55	Cs	caesium 133	87	Ļ	francium -
	IIV V VI	Group	1	II	III	1	II	II	II	III	II	II	II	II	II	II	1   1   1   1   1   1   1   1   1   1	1	1	1

71	P	Iutetium	175	103	۲	lawrencium	ı
70	Хp	ytterbium	173	102	8	nobelium	ı
69	Tm	thulium	169	101	Md	mendelevium	1
89	щ	erbium	167	100	Fm	fermium	1
29	운	holmium	165	66	Es	einsteinium	ı
99	۵	dysprosium	163	86	ర	californium	I
65	Д	terbium	159	26	BK	berkelium	1
64	Gd	gadolinium	157	96	Cm	curium	1
63	En	europium	152	96	Am	americium	I
62	Sm	samarium	150	94	Pn	plutonium	ı
61	Pm	promethium	ı	93	ď	neptunium	ı
09	PZ	neodymium	144	92	$\supset$	uranium	238
59	Ą						
28	Ce	cerium	140	06	Ļ	thorium	232
22	Га	lanthanum	139	88	Ac	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).