



## **Cambridge Assessment International Education**

Cambridge International General Certificate of Secondary Education (9–1)

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0971/42

Paper 4 Theory (Extended)

October/November 2019

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

## **READ THESE INSTRUCTIONS FIRST**

Write your centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 12.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

This document consists of 12 printed pages.



1 The Periodic Table is very useful to chemists.

Refer only to elements with atomic numbers 1 to 36 in the Periodic Table provided when answering **Question 1**.


(a)	Use	information from the Periodic Table provided to identify <b>one</b> element which:	
	(i)	has atoms with exactly 9 protons	[1]
	(ii)	has atoms with 0 neutrons	[1]
(	iii)	has atoms with exactly 23 electrons	[1]
(	iv)	has atoms with an electronic structure of 2,8,6	[1]
	(v)	forms ions with a charge of 3– containing 18 electrons	[1]
(	vi)	forms ions with a charge of 2+ containing 10 electrons	[1]
(v	/ii)	has a relative atomic mass that shows it has at least two isotopes	[1]
(b)	Stat	te which metal in the first 36 elements:	
	(i)	is the Group I element which reacts most vigorously with water	[1]
	(ii)	reacts with air to form lime.	[1]
	One	e element in the first 36 elements is used as the fuel in a fuel cell.  Name this element.	
	(-)		[1]
(	(ii)	Write the overall chemical equation for the reaction which occurs when the element (c)(i) reacts in a fuel cell.	
			[2]
		[Total:	12]

٦	The gases Ar, CO <sub>2</sub> , N <sub>2</sub> and O <sub>2</sub> are in clean, dry air.	
(	CO, NO, NO <sub>2</sub> and SO <sub>2</sub> are gases commonly found in polluted air.	
(	a) What percentage of clean, dry air is N <sub>2</sub> ?	
	Give your answer to the nearest whole number.	
	%	, [1]
(	Name the process used to separate $O_2$ from clean, dry air.	[2]
(	c) State <b>one</b> major adverse effect of the pollutant SO <sub>2</sub> .	· [∠]
		[1]
(	d) NO and NO <sub>2</sub> are produced in car engines.	
	Describe how oxides of nitrogen form in a car engine.	
		[2]
(	Many cars have catalytic converters in their exhaust systems. In a catalytic converter, mo the CO and NO formed in a car engine is changed into less harmful products.	st of
	Identify these products and state the metal catalyst used.	
	products	
	catalyst	 [3]
(	f) CO is formed from the incomplete combustion of fossil fuels such as methane.	
	Write a chemical equation to show the incomplete combustion of methane.	
		. [2]

2

(g) The  ${\rm CO_2}$  in air is part of the carbon cycle.

The scheme shows a simple representation of part of the carbon cycle.



	in green plants	
(i)	State the scientific terms for each of process <b>A</b> and process <b>B</b> .	
	A	
	В	 [2]
(ii)	Plants convert glucose into complex carbohydrates.	
	A unit of glucose can be represented as HO—OH.	
	Complete the diagram to show the complex carbohydrate formed from <b>three</b> units glucose. Show all of the atoms and all of the bonds in the linkages.	of
		[2]
(iii)	Complex carbohydrates break down to form simple sugars.	
	State <b>two</b> ways that complex carbohydrates can be broken down into simple sugars.	
	1	
	2	 [2]
		[4]
(iv)	Name a suitable technique for separating and identifying the individual sugars form when complex carbohydrates are broken down.	ed
		[1]

[Total: 18]

© UCLES 2019 0971/42/O/N/19

3	Am	mon	ia is an important chemical.
	(a)	Am	monia is manufactured by the Haber process. The reaction is reversible.
		(i)	What is the sign for a reversible reaction?
			[1]
		(ii)	State the essential conditions for the manufacture of ammonia by the Haber process starting from hydrogen and nitrogen. Include a chemical equation to show the reaction which occurs.
			[5]
	(	(iii)	Name <b>one</b> raw material which is a source of the hydrogen used in the Haber process.
			[1]
	(b)	Am	monia is a base and reacts with sulfuric acid to form the salt, ammonium sulfate.
		(i)	What is meant by the term base?
			[1]
		(ii)	Name the industrial process used to manufacture sulfuric acid.
		(iii)	Write a chemical equation for the reaction between ammonia and sulfuric acid.
	,	(**** <i>)</i>	[2]
			[-]

(i) Name the green precipitate.	ſhis
	[1]
(ii) Suggest why the green precipitate turns red-brown at the surface.	
	[2]
(iii) State what happens when an excess of aqueous ammonia is added to the green precipit	ate.
	[1]

(d) Ammonia reacts with oxygen as show	(d)	Ammonia	reacts with	oxygen	as	show
--	-----	---------	-------------	--------	----	------

$$4NH_3(g) + 5O_2(g) \rightarrow 4NO(g) + 6H_2O(g)$$

(i)	Calculate the volume of oxygen at room temperature and pressure, in dm³, that reacts with
	4.80 dm <sup>3</sup> of ammonia.

(ii) The chemical equation for the reaction can be represented as shown.

4 H—N—H + 5 O=O 
$$\rightarrow$$
 4 N=O + 6 H—O—H H

Use the bond energies in the table to calculate the energy change, in kJ/mol, which occurs when **one** mole of  $NH_3$  reacts.

bond	N–H	O=O	N=O	О–Н
bond energy in kJ/mol	391	498	587	464

Energy needed to break bonds.

.....kJ

Energy released when bonds are formed.

..... kJ

• Energy change when **one** mole of NH<sub>3</sub> reacts.


[Total: 22]

4	Many	substances	conduct	electricity.
	IVICIII	Cabotalloco	COLIGACE	Old Oth Tolky

(a)	Identify all the particles responsible for the passage of electricity in:

•	• graphite	
---	------------	--

- molten copper(II) bromide. .....
  [4]
- **(b)** A student used the following apparatus to electrolyse concentrated aqueous sodium chloride using inert electrodes.




Suggest the name of a metal which could be used as the inert electrodes.
[1]
Name the gas formed at the positive electrode.
[1]
Write an ionic half-equation for the reaction occurring at the negative electrode. Include state symbols.
[3]
How, if at all, does the pH of the solution change during the electrolysis? Explain your answer.

© UCLES 2019 0971/42/O/N/19

(c) A student used the following electrochemical cell.

The reading on the voltmeter was +1.10 V.



(i)	Draw an arrow on the diagram to show the direction of electron flow.	[1]
(ii)	Suggest the change, if any, in the voltmeter reading if the zinc electrode was replaced van iron electrode. Explain your answer.	vith
		[2]
(iii)	The zinc electrode was replaced with a silver electrode. The reading on the voltmeter v –0.46 V.	vas
	Suggest why the sign of the voltmeter reading became negative.	
		[1]

[Total: 16]

- **5** Methanol, CH<sub>3</sub>OH, is a member of the homologous series of alcohols.
  - (a) Methanol can be made from methane in a two-step process.
    - **step 1** Methane is reacted with chlorine gas to produce chloromethane, CH<sub>3</sub>C*l*.
    - **step 2** CH<sub>3</sub>C*l* is reacted with sodium hydroxide to produce CH<sub>3</sub>OH and one other product.
    - (i) What conditions are needed in step 1?

F	F 4	-
	17	
	4 1	

(ii) Write the chemical equation for the reaction which occurs in step 1.


·		-
	11	- 1
		- 1

(iii) State the type of organic reaction occurring in **step 1**.

(iv) Complete the chemical equation for step 2.

$$CH_3Cl + NaOH \rightarrow CH_3OH + \dots$$
 [1]

**(b)** Draw a dot-and-cross diagram to show the electron arrangement in a molecule of methanol. Show outer shell electrons only.



[2]

c) Me	thanol reacts with propanoic acid to form an ester with a molecular formula $C_4H_8O_2$ .
(i)	Name the ester formed when methanol reacts with propanoic acid.
	[1]
(ii)	Name one other substance formed when methanol reacts with propanoic acid.
	[1]
(iii)	Draw the structure of an ester which is a structural isomer of the ester named in <b>(c)(i)</b> . Show all of the atoms and all of the bonds.
	[3]
(iv)	State the conditions needed to form an ester from a carboxylic acid and an alcohol.
	[1]
	[Total: 12]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

The Periodic Table of Elements

	<b>=</b>	5	e H	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton 84	25	Xe	xenon	98	R	radon			
	<b>=</b>				6	ш	fluorine 19	17	Cl	chlorine 35.5	35	B	bromine 80	53	н	iodine 127	85	¥	astatine -			
-	<b>=</b>																			116		ermorium -
:	>																					≧
	<u> </u>									<u> </u>										114	Εl	erovium -
	=																					
										a ———										112	C	pernicium
																						darmstadtium roe
Group																_						_
			_	nago																		
			_	hydro 1																		
								1			25	Mn	mangane 55	43								
					_	pol	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
				Key	atomic numbe	mic sym	name ative atomic m				23	>	vanadium 51	41	g	niobium 93	73	٦	tantalum 181	105	Op	dubnium —
						ato	rela				22	ı=	titanium 48	40	Zr	zirconium 91	72	士	hafnium 178	104	꿆	rutherfordium -
								•			21	Sc	scandium 45	39	>	yttrium 89	57–71	lanthanoids		89–103	actinoids	
	=				4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ഗ്	strontium 88	56	Ba	barium 137	88	Ra	radium -
	_				8	:=	lithium 7	11	Na	sodium 23	19	¥	potassium 39	37	Rb	rubidium 85	55	Cs	caesium 133	87	Ļ	francium —
			Group  III IV V V VII	Group  III IV V V VII	1	III	II	III	II	II	II	II	II	II	II	II	II	II	1	1	1	II

71	Γn	Iutetium	175	103	۲	lawrencium	I
	Υp					_	
69	H	thulium	169	101	Md	mendelevium	1
89	ш	erbinm	167	100	Fm	ferminm	1
29	웃	holmium	165	66	Es	einsteinium	I
99	۵	dysprosium	163	86	ర్	californium	1
65	q	terbium	159	97	BK	berkelium	1
64	Gd	gadolinium	157	96	Cm	curium	1
63	En	europium	152	92	Am	americium	1
62	Sm	samarium	150	94	Pu	plutonium	I
61	Pm	promethium	I	93	d d	neptunium	ſ
09	pN	neodymium	144	92	$\supset$	uranium	238
59	Ā	praseodymium	141	91	Ра	protactinium	231
28	Ce	cerium	140	06	드	thorium	232
22	Гa	lanthanum	139	89	Ac	actinium	I

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).