

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**  
International General Certificate of Secondary Education

**MARK SCHEME for the October/November 2013 series**

<b>0444 MATHEMATICS (US)</b>	
<b>0444/23</b>	Paper 2 (Extended), maximum raw mark 70

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

<b>Page 2</b>	<b>Mark Scheme</b>	<b>Syllabus</b>
	<b>IGCSE – October/November 2013</b>	<b>0444</b>

**Abbreviations**

- cao correct answer only
- cso correct solution only
- dep dependent
- ft follow through after error
- isw ignore subsequent working
- oe or equivalent
- SC Special Case
- www without wrong working
- art anything rounding to
- soi seen or implied

	<b>Answers</b>	<b>Mark</b>	<b>Part Marks</b>
<b>1</b>	39	<b>2</b>	<b>M1</b> for $52 \times 45 \div 60$ oe
<b>2</b>	Any two of (20, 8) (-4, 0) (12, 24)	<b>2</b>	<b>B1</b> for one correct
<b>3</b>	-8	<b>2</b>	<b>M1</b> for $2x = -16$ or $\frac{1}{2} + x = -7.5$
<b>4</b>	64	<b>2</b>	<b>M1</b> for (their $(5 - 1)^3$ )
<b>5</b>	[domain] $0 \leq x \leq 3$ [range] $2$	<b>1</b> <b>1</b>	
<b>6 (a)</b>	600 000	<b>1</b>	
<b>(b)</b>	90	<b>2</b>	<b>M1</b> for $\div 1000 \times 60 \times 60$
<b>7</b>	30	<b>3</b>	<b>M2</b> for $24 \div 0.8$ or <b>M1</b> for recognition of $80\% = 24$
<b>8</b>	5	<b>3</b>	<b>M2</b> for $(x - 5)(x - 1)$ or <b>M1</b> for evidence of a factorisation which gives the correct coefficient of $x$ or positive prime constant term e.g. $(x - 7)(x + 1)$ , $(x - 4)(x - 2)$ , $(x - 3)(x - 1)$
<b>9</b>	1600	<b>3</b>	<b>M1</b> for $m = kx^3$ <b>A1</b> $k = 25$ or <b>M2</b> for $200 \times \left(\frac{4}{2}\right)^3$
<b>10 (a)</b>	$a^2 + 2ab + b^2$ final answer	<b>2</b>	<b>B1</b> for $a^2$ [+] $ab$ [+] $ab$ [+] $b^2$ seen
<b>(b)</b>	22	<b>1</b>	
<b>11</b>	12	<b>3</b>	<b>M2</b> for $\sqrt{15^2 - 9^2}$ or <b>M1</b> for $AB^2 + 9^2 = 15^2$ oe

Page 3	Mark Scheme	Syllabus
	IGCSE – October/November 2013	0444

12 (a)	[amplitude] 2 [period] 360	1 1	
(b)	$4 \sin x$	1	
13 (a)	2	1	
(b)	Accurate bisector of <b>either</b> side of rectangle	2	<b>B1</b> for correct ruled line (cross two sides) <b>B1</b> for 2 pairs of correct arcs
14 (a)	$4.8 \times 10^6$	2	<b>B1</b> for 4 800 000
(b)	$9.3 \times 10^7$	2	<b>B1</b> for 93 000 000 or $93 \times 10^6$ or $0.93 \times 10^8$ oe
15 (a)	24	2	<b>M1</b> for $MOC = 48$
(b)	24	2	<b>M1</b> for $ACM = 66$ or <b>B1</b> for 48 – <i>their (a)</i>
16 (a)	$8q^{-1}$ or $\frac{8}{q}$	2	<b>B1</b> for $8q^k$ or $kq^{-1}$
(b)	$\frac{1}{5}$ or 0.2	2	<b>M1</b> for $5^{-2}$ , $\frac{1}{5^2}$ or [0].04seen oe
17 (a)	triangle at (0, 2) (0, 4) and (-1, 2)	2	<b>SC1</b> for rotation $90^\circ$ clockwise about (0, 1) or any other rotation $90^\circ$ anticlockwise
(b)	stretch x-axis invariant [factor] 2	1 1 1	
18	[c = ] 6 [d = ] 9	4	accept any correct method e.g. <b>M1</b> for $\frac{30}{360} \times \pi \times 6^2$ [ $\times 2$ ] <b>A1</b> for $6\pi$ or 6  <b>M1</b> for $\frac{1}{2} \times 6^2 \times \sin 120$ or <b>B1</b> for $\sin 120 = \frac{\sqrt{3}}{2}$
19 (a)	19 – 19.1	1	
(b)	3	2	<b>M1</b> for 47 seen
(c)	4.9 to 5.7	2	<b>B1</b> for [UQ] 21.7 to 22.2 and [LQ] 16.5 to 16.8
(d)	$\frac{45}{50}$ oe	2	<b>B1</b> for 45 seen or <b>SC1</b> for $\frac{5}{50}$ isw

<b>Page 4</b>	<b>Mark Scheme</b>	<b>Syllabus</b>
	<b>IGCSE – October/November 2013</b>	<b>0444</b>

<b>20 (a)</b>	75	<b>2</b>	<b>B1</b> for $[g(6)=] 36$
<b>(b)</b>	3.5    -6.5	<b>3</b>	<b>M1</b> for $(2x + 3)^2 = 100$ <b>M1</b> for $2x + 3 = [\pm]10$  if 0 scored <b>SC1</b> for one correct value as answer
<b>(c)</b>	$\frac{x-3}{2}$ oe final answer	<b>2</b>	<b>M1</b> for $x = 2y + 3$ or $y - 3 = 2x$ or $\frac{y}{2} = x + \frac{3}{2}$
<b>(d)</b>	5	<b>1</b>	