

**MARK SCHEME for the October/November 2011 question paper  
for the guidance of teachers**

**0606 ADDITIONAL MATHEMATICS**

**0606/12**

Paper 1, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2011	0606	

## Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
  - A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
  - B Accuracy mark for a correct result or statement independent of method marks.
- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep\*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
  - The symbol  $\surd$  implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
  - Note: B2 or A2 means that the candidate can earn 2 or 0.  
B2, 1, 0 means that the candidate can earn anything from 0 to 2.

<b>Page 3</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE – October/November 2011</b>	<b>0606</b>	

The following abbreviations may be used in a mark scheme or used on the scripts:

- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)

### Penalties

- MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through  $\sqrt{\quad}$ ” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy.
- OW –1,2 This is deducted from A or B marks when essential working is omitted.
- PA –1 This is deducted from A or B marks in the case of premature approximation.
- S –1 Occasionally used for persistent slackness – usually discussed at a meeting.
- EX –1 Applied to A or B marks when extra solutions are offered to a particular equation. Again, this is usually discussed at the meeting.

<b>Page 4</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>
	<b>IGCSE – October/November 2011</b>	<b>0606</b>

<p><b>1</b></p> $\frac{1}{\frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta}}$ $= \frac{\sin \theta \cos \theta}{\sin^2 \theta + \cos^2 \theta}$ $= \sin \theta \cos \theta$ <p><b>OR</b></p> $\frac{\tan \theta}{\tan^2 \theta + 1} \text{ or } \frac{\cot \theta}{\cot^2 \theta + 1}$ $= \frac{\tan \theta}{\sec^2 \theta} \text{ or } \frac{\cot \theta}{\operatorname{cosec}^2 \theta}$ $= \sin \theta \cos \theta$	<p>M1</p> <p>M1</p> <p>A1 [3]</p> <p>M1</p> <p>M1</p> <p>A1</p>	<p>M1 for adding fractions in terms of sin/cos/tan/cot correctly</p> <p>M1 for use of correct identity</p> <p>A1 for correct solution only</p> <p>M1 for adding fractions in terms of tan/cot correctly</p> <p>M1 for use of correct identity</p> <p>A1 for correct solution only</p>
<p><b>2</b></p> $(2y+1)^2 + y^2 = 29$ <p>(or <math>5x^2 - 2x - 115 = 0</math>)</p> <p>leading to <math>5y^2 + 4y - 28 = 0</math></p> <p>(or <math>x^2 + \left(\frac{x-1}{2}\right)^2 = 29</math>)</p> $x = -\frac{23}{5}, y = -\frac{14}{5} \text{ and}$ $x = 5, y = 2$ <p>(5, 2) spotted gets B1</p>	<p>M1</p> <p>DM1</p> <p>DM1</p> <p>A1</p> <p>A1</p> <p>[5]</p>	<p>M1 for attempt to get an equation in terms of one variable only</p> <p>DM1 for obtaining a 3 term quadratic equation</p> <p>DM1 for attempt to solve quadratic equation</p> <p>A1 for a pair of values</p>

Page 5	Mark Scheme: Teachers' version	Syllabus
	IGCSE – October/November 2011	0606

<p>3 (i) <math>\frac{1}{\log_2 x}</math> or <math>\frac{\log_2 2}{\log_2 x}</math></p> <p>(ii) <math>u^2 - 3u + 2 = 0</math>  <math>(u - 1)(u - 2) = 0</math></p> <p><math>u = 1, 2</math></p> <p><math>x = 2</math> and <math>x = 4</math></p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>[5]</p>	<p>M1 for a correct attempt to obtain and solve a quadratic equation in terms of <math>u</math> or <math>\log_2 x</math></p> <p>A1 for <math>u = 1, 2</math></p> <p>M1 for attempt to solve an equation of the form <math>\log_2 x = k</math> leading to <math>x = 2^k</math></p> <p>A1 for both</p>
<p>4 When <math>x = 2, y = 9</math></p> <p><math>\frac{dy}{dx} = \frac{2}{3} \cdot 6x \cdot (3x^2 + 15)^{-\frac{1}{3}}</math></p> <p>when <math>x = 2, \frac{dy}{dx} = \frac{8}{3}</math></p> <p><math>\therefore</math> grad of normal = <math>-\frac{3}{8}</math></p> <p>normal : <math>y - 9 = -\frac{3}{8}(x - 2)</math></p> <p><math>8y + 3x = 78</math></p>	<p>B1</p> <p>B1, B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[6]</p>	<p>B1 for <math>y = 9</math></p> <p>B1 for <math>\frac{2}{3} \cdot 6x</math>, B1 for <math>(3x^2 + 15)^{-\frac{1}{3}}</math></p> <p>M1 for use of <math>m_1 m_2 = -1</math></p> <p>M1 for attempt to find equation of normal, must be using gradient of a perpendicular line</p> <p>A1 allow unsimplified</p>
<p>5 (i) <math>y^2 = m2^x + c</math></p> <p><math>81 = 80 + c</math>  <math>c = 1</math></p> <p><math>y^2 = 5(2^x) + 1</math></p> <p>(ii) <math>36 = 5(2^x) + 1</math></p> <p>leading to <math>7 = 2^x</math></p> <p><math>x = 2.81</math></p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>DM1</p> <p>A1</p> <p>[6]</p>	<p>M1 for use of straight line equation as given</p> <p>M1 for use of (16, 81)</p> <p>A1 – do not allow if subsequent incorrect work is seen</p> <p>M1 substitution of <math>y = 6</math> into their equation in terms of <math>y^2</math> and <math>2^x</math></p> <p>DM1 for correct solution of equation of the form <math>2^x = k</math></p>

Page 6	Mark Scheme: Teachers' version	Syllabus
	IGCSE – October/November 2011	0606

<p>6 (i)</p> $243 + 405x + 270x^2 + 90x^3 + 15x^4 + x^5$ $243 - 405x + 270x^2 - 90x^3 + 15x^4 - x^5$ $486 + 540x^2 + 30x^4$ <p>(ii) <math>30y^2 + 540y - 600 = 0</math></p> <p><math>y = 1.05</math></p> <p>leading to <math>x = \pm 1.02</math></p>	<p>B1 B1 B1 B1</p> <p>M1 DM1</p> <p>A1 A1</p> <p>[8]</p>	<p>Assuming correct terms in <math>x</math> B1 for 486 B1 for 540 B1 for 30 B1 for all terms correct and no extra terms</p> <p>M1 for attempt to obtain a 3 term quadratic equation. DM1 for correct attempt at solution of quadratic</p> <p>A1 need both solutions A1 need both solutions</p>
<p>7 (i) <math>16x^{-\frac{1}{2}} - 8 + x^{\frac{1}{2}}</math></p> <p>(ii) <math>y = 32x^{\frac{1}{2}} - 8x + \frac{2}{3}x^{\frac{3}{2}} (+c)</math></p> <p>When <math>x = 9</math> and <math>y = 30</math>, <math>c = -12</math></p>	<p>B1, B1, B1</p> <p>M1 A2, 1, 0</p> <p>M1 A1</p> <p>[8]</p>	<p>B1 for each correct term</p> <p>M1 for attempt to integrate a 3 term expression -1 for each error</p> <p>M1 for attempt to find <math>c</math>, must have attempted integration A1 for <math>c = -12</math></p>
<p>8 (i) <math>M(2, 4)</math></p> <p><math>\text{Grad } AB = \frac{-2}{6}</math>, <math>\perp</math> grad = 3</p> <p><math>CD: y - 4 = 3(x - 2)</math></p> <p>(ii) <math>C(-2, 2)</math> <math>D(10, -7)</math></p> <p>(iii) Area = 37.5</p> <p>Special case: (ii) allow <math>D(-6, 5)</math> (iii) area 12.5</p>	<p>B1</p> <p>M1</p> <p>DM1 A1</p> <p>B1 B1, B1</p> <p>M1, A1</p> <p>[9]</p> <p>B1, B1 M1, A1</p>	<p>Allow in (ii)</p> <p>M1 for attempt to find gradient of perpendicular</p> <p>DM1 for straight line equation</p> <p>B1 for <math>y</math> coordinate of <math>A</math> B1 for each</p> <p>M1 for a valid attempt at area</p>

Page 7	Mark Scheme: Teachers' version	Syllabus
	IGCSE – October/November 2011	0606

<p>9 (i) <math>\frac{dy}{dx} = x.4 \cos 4x + \sin 4x</math></p> <p>(ii) <math>I = \left(\frac{1}{4}\right) \left[ x \sin 4x - \int \sin 4x \, dx \right]</math></p> $= \left(\frac{1}{4}\right) \left[ x \sin 4x - \left(-\frac{1}{4} \cos 4x\right) \right]$ <p>For definite integral</p> $\left(\frac{1}{4}\right) \left[ x \sin 4x - \left(-\frac{1}{4} \cos 4x\right) \right]_0^{\frac{\pi}{8}}$ $= \frac{\pi}{32} - \frac{1}{16}, \text{ or } 0.0357$	<p>M1 B1, A1</p> <p>DM1</p> <p>A1, A1 B1</p> <p>M1</p> <p>A1 [9]</p>	<p>M1 for differentiation of a product B1 for <math>4 \cos 4x</math>, A1 all else correct</p> <p>DM1 for realising integration is form reverse process of (i) – do not need <math>\left(\frac{1}{4}\right)</math> until last A1</p> <p>A1 for <math>x \sin 4x</math>, A1 for <math>\int \sin 4x \, dx</math> B1 for <math>-\frac{1}{4} \cos 4x</math></p> <p>M1 for correct application of limits</p>
<p>10 (i) <math>2 \tan^2 x + 2 = 5 \tan x + 5</math></p> $2 \tan^2 x - 5 \tan x - 3 = 0$ $\tan x = -\frac{1}{2}, \tan x = 3$ $x = 153.4^\circ, 333.4^\circ \text{ and } 71.6^\circ, 251.6^\circ$ <p>(ii) <math>\sin\left(0.5y + \frac{\pi}{3}\right) = \frac{1}{\sqrt{2}}</math></p> $0.5y + \frac{\pi}{3} = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{9\pi}{4}$ <p>leading to <math>y = \frac{5\pi}{6}, \frac{23\pi}{6}</math></p>	<p>M1</p> <p>M1</p> <p>M1</p> <p>A1, A1</p> <p>M1</p> <p>M1 M1</p> <p>A1, A1 [10]</p>	<p>M1 for use of correct identity</p> <p>M1 for solution of 3 term quadratic equation</p> <p>M1 for attempt to solve <math>\tan x = k</math> from a 3 term quadratic equation</p> <p>A1 for any pair</p> <p>M1 for dealing with <math>\sqrt{2}</math> correctly</p> <p>M1 for correct order of operations M1 for correct order of operations and attempt to get a solution in the range</p> <p>Allow decimal equivalents 2.62 and 12.0</p>

Page 8	Mark Scheme: Teachers' version	Syllabus
	IGCSE – October/November 2011	0606

<p><b>11 EITHER</b></p> <p>(i) <math>A = 4</math></p> <p>(ii) <math>\frac{dy}{dx} = e^{-x}(-2A \sin 2x + 2B \cos 2x) - e^{-x}(A \cos 2x + B \sin 2x)</math></p> <p>when <math>x = 0, 6 = 2B - A, B = 5</math> (verification acceptable)</p> <p>(iii) when <math>\frac{dy}{dx} = 0,</math></p> <p><math>e^{-x}(p \cos 2x - q \sin 2x) = 0</math></p> <p>leading to <math>\tan 2x = \frac{p}{q} \left( = \frac{6}{13} \right)</math></p> <p><math>x = 0.216</math></p>	<p>B1</p> <p>M1 B1</p> <p>B1</p> <p>M1 A1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>M1 A1</p> <p>[11]</p>	<p>M1 for differentiation of a product B1 for <math>(-2A \sin 2x + 2B \cos 2x)e^{-x}</math></p> <p>B1 for <math>-e^{-x}(A \cos 2x + B \sin 2x)</math></p> <p>M1 for substitution to find <math>B</math></p> <p>M1 for their <math>\frac{dy}{dx} = 0</math></p> <p>M1 for attempt to simplify</p> <p>M1 for attempt to obtain <math>\tan 2x = \frac{p}{q}</math></p> <p>M1 for attempt to solve <math>\tan 2x = k</math></p>
<p><b>11 OR</b></p> <p>(i) <math>\frac{dy}{dx} = \frac{(x^2 - 1) \frac{2x}{(x^2 - 1)} - 2x \ln(x^2 - 1)}{(x^2 - 1)^2}</math></p> <p>Rearranging to get <math>k = 2</math></p> <p>(ii) <math>\partial y = \frac{dy}{dx} p,</math></p> <p>leading to <math>\partial y = -0.108p</math></p> <p>(iii) when <math>\frac{dy}{dx} = 0, 1 - \ln(x^2 - 1) = 0</math></p> <p><math>\ln(x^2 - 1) = 1</math></p> <p><math>x^2 - 1 = e</math> or 2.72</p> <p>leading to <math>x = \sqrt{1 + e}</math></p> <p><math>y = \frac{1}{e}</math></p>	<p>M1 B1</p> <p>A1 A1</p> <p>M1</p> <p>√A1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>[11]</p>	<p>M1 for differentiation of a quotient B1 <math>\frac{2x}{(x^2 - 1)}</math></p> <p>A1 for all else correct A1 for rearrangement to get <math>k = 2</math></p> <p>M1 for substitution of <math>x = \sqrt{5}</math> and correct method</p> <p>√A1 on their <math>k</math></p> <p>M1 for <math>\frac{dy}{dx} = 0</math> and attempt to simplify</p> <p>A1 for <math>\ln(x^2 - 1) = 1</math></p> <p>A1 for <math>x^2 - 1 = e</math> or 2.72</p>