


Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		



PHYSICAL SCIENCE

0652/41

Paper 4 Theory (Extended)

October/November 2021

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

This document has 24 pages. Any blank pages are indicated.

DC (RW/CB) 212508/5 © UCLES 2021

[Turn over

1 A skydiver of mass 85 kg jumps out of an aircraft and initially falls freely under gravity.




Fig. 1.1

(a) (i) Calculate the force due to gravity acting on the skydiver. [g = 10 N/kg]

			force =		N [2]
	(ii)	Deduce the initial acceleration	of the skydiver.		
			acceleration =		m/s^2 [1]
(b) The acceleration of the skydiver decreases as she falls, even though she does not parachute.					
	Ехр	plain why her acceleration decrea	ases.		
					[2]

(c) The graph in Fig. 1.2 shows how the speed of the skydiver changes as she falls before opening her parachute.




Fig. 1.2

(i) At time *t* the skydiver is falling with constant speed.

State the term used to describe this constant speed.

[1]

(ii) State the magnitude of the resultant force on the skydiver when her speed is constant.

magnitude of resultant force =[1]

[Total: 7]

2 The formulae of some substances are shown in Fig. 2.1.

CaO	CH₃CH₂OH	CH₃OH	C ₂ H ₄	
C_2H_6	Cl ₂	СО	CO ₂	
	Fig. 2	.1		

(a) Answer the following questions by selecting from the substances shown in Fig. 2.1.

Each substance may be used once, more than once or not at all.

State the formula for a substance which:

(i)	is a diatomic gas at room temperature	
		[1]
(ii)	is used to reduce emissions of sulfur dioxide	
		[1]
(iii)	is a solvent produced during a reaction involving yeast and glucose	
		[1]
(iv)	turns aqueous bromine colourless	
		[1]
(v)	is a neutral oxide	
(vi)	is a gas produced during the catalytic removal of nitrogen monoxide from exha emissions.	ust
		[1]

(b) CH₃OH is a covalent molecule.

Complete the dot-and-cross diagram in Fig. 2.2 to show the bonding in one molecule of $\mathrm{CH_3OH}.$

Show the outer electrons only.




Fig. 2.2

[2]

[Total: 8]

3 Fig. 3.1 shows a mechanic using a wrench to tighten a nut on the wheel of a car.




Fig. 3.1

The nut acts as a pivot.

Fig. 3.2 shows the front view of the wrench.




Fig. 3.2

- (a) The force applied by the mechanic is a steady 240 N. The distance *d* of the force from the centre of the nut (pivot) is 0.45 m.
 - (i) Calculate the moment produced about the centre of the nut (pivot).

moment =Nm [2]

(ii) Describe how the moment about the nut (pivot) can be increased without changing the magnitude of the force.

(b)	The 240N force rotates the wrench through 360°. The distance moved by the force is 2.8 m.
	Calculate the work done by the force. State the unit.

work done =		unit	[3]
-------------	--	------	-----

[Total: 6]

BLANK PAGE

	aidiii	chloride is an ionic compound.							
(a)	(a) In an experiment, molten sodium chloride is electrolysed.								
	(i)	Explain the meaning of <i>electrolysis</i> .							
		[2]							
	(ii)	Predict the products formed at each electrode during the electrolysis of molten sodium chloride.							
		anode							
		cathode							
		[2]							
	(iii)	Describe what happens, in terms of electrons, to the cations in sodium chloride during electrolysis.							
		[1]							
	(iv)	The sodium chloride in this experiment is molten. This is shown by using the state symbol (I) in the formula $NaCl(I)$.							
		Complete the formula to show sodium chloride in the other state in which it can be							
		electrolysed.							
		electrolysed. NaCl() [1]							
(b)	Des								
(b)		NaCl() [1] scribe two features of the structure of ionic compounds such as sodium chloride.							
(b)		NaCl() Scribe two features of the structure of ionic compounds such as sodium chloride.							
(b)	1	NaCl() Scribe two features of the structure of ionic compounds such as sodium chloride.							
(b)	1	NaCl() Scribe two features of the structure of ionic compounds such as sodium chloride.							
(b)	1	NaCl() Scribe two features of the structure of ionic compounds such as sodium chloride.							

4

5 A rectangular block of wood vibrates to produce waves in a ripple tank. The wavefronts travel towards a narrow gap in a barrier, as shown in Fig. 5.1.




Fig. 5.1

- (a) (i) On Fig. 5.1 draw a double-headed arrow (↔) to show one wavelength of the waves. [1]
 (ii) Name the process that occurs as the waves go through the gap in the barrier.
 [1]
 (iii) On Fig. 5.1, draw three wavefronts after they have passed through the gap. [3]
- (b) (i) The vibrating block of wood is 28 cm from the barrier.

Each wavefront takes 0.80s to travel from the block to the barrier.

Calculate the speed of the wave.

speed =cm/s [2]

	(ii)		avelength		waves	is 5.6	cm. Use	your	answei	r to	(b)(i)	to	calculate	the
						freq	uency =						H	z [2]
													[Tota	al: 9]
(a)		mplete ī lium.	Table 6.1 t	to show	the nu	mber of	protons	s, neut	rons an	d ele	ectron	ıs ir	n one ato	m of
						Table (6.1							
			number	of proto	ons	number	of neut	rons	numbe	er of	electr	ons	3	
		²³ Na												
			l											[2]
(b)	The	ere are s	several diff	ferent iso	otopes	of sodiu	ım.							
	Explain why isotopes of an element have the same properties.													
														. [1]
													[Tota	al: 3]

6

7 A metal sphere is suspended by a plastic thread.

A conducting wire is attached to the positive terminal of a power supply and placed near to the sphere as shown in Fig. 7.1.




Fig. 7.1

(a) The electromotive force (e.m.f.) of the high voltage power supply is 5000 V.

A safety resistor of resistance $7.5\,M\Omega$ is connected between the conducting wire and the positive terminal of the power supply.

Explain why this resistor enables the power supply to be used safely.							
					[1]		

(b) The conducting wire is connected to the metal sphere. There is an average current of 0.67 mA for a time of 0.25 ms.

Calculate the charge transferred to the metal sphere.

(C)	A small charged sphere is held near the large metal sphere after the conducting wire is removed.
	The small sphere is repelled by the electric field of the large sphere.
	Explain what is meant by an electric field.
	[1]
(d)	Suggest why the large metal sphere is suspended by a plastic thread rather than a copper wire.
	[2]
	[Total: 6]

Car	bon is an element in Group IV and Period 2 of the Periodic Table.
(a)	The position of carbon in the Periodic Table gives two pieces of information about its electronic structure.
	State these two pieces of information.
	1
	2[2]
(b)	Graphite is a form of carbon.
	Explain, in terms of structure, why graphite can be used as a lubricant.
	[2]
(c)	Carbon is used in the extraction of iron from its ore in a blast furnace.
	Carbon reacts with oxygen in the blast furnace to form carbon dioxide.
	$C + O_2 \rightarrow CO_2$
	Write two other equations for essential reactions that occur in the blast furnace to extract iron from its ore.
	1
	2[2]

© UCLES 2021 0652/41/O/N/21

8

4	/ _I \	Zinc blende	:	- f :	14	70
1	aı	/ Inc hiende	is an ore	OT ZINC	it contains	/ns
л	M.		13 411 010		it contains	

Before zinc metal is extracted, its ore is crushed and heated in air.

2ZnS +
$$3O_2 \rightarrow 2ZnO + 2SO_2$$

Calculate the mass of ZnO that is produced from 6.5 tonnes of ZnS.

 $[A_r: Zn, 65; S, 32; O, 16; and 1 tonne = 1000 kg]$

Give your answer to two significant figures.

		mass of ZnO =	 tonnes	[3]

(e)	Ste	el alloys contain iron. Steel is stronger and harder than iron.	
	(i)	State the meaning of alloy.	
			[1]
	(ii)	Fig. 8.1 is a diagram of a model of the atoms in pure iron metal.	




Fig. 8.1

Draw a diagram in the box to show a model of the atoms in an alloy of iron.



[1]

[Total: 11]

9 Fig. 9.1 is a circuit diagram containing a battery of electromotive force (e.m.f.) 6.0 V, a resistor of resistance 2.2Ω , an ammeter, a switch and a lamp.




Fig. 9.1

When the switch is closed the reading on the ammeter is 1.2A.

(a) (i) Calculate the total resistance in the circuit.

total resistance =	Ω	[2]
of the lamp		

(ii) Calculate the resistance of the lamp.

resistance of the lamp =	Ω	[1]]
--------------------------	----------	-----	---

(b) The circuit is reconnected so that the lamp and the resistor are connected in parallel with each other.

Explain why the lamp now shines more brightly than when connected in the circuit shown in Fig. 9.1.

 	• • •	 	 	 	٠.	 	 	 	 	 	 •••	 	 	 														
 		 	 	 		 	 	 	 	 	 	 	 		[2]													

[Total: 5]


10	(a)	Cal	cium carbonate reacts with dilute hydrochloric acid.
		Def	ine, by referring to proton transfer, what is meant by acid.
			[1]
	(b)		e of the products of the reaction between calcium carbonate and dilute hydrochloric acid is bon dioxide gas.
		(i)	Write the symbol equation for this reaction.
			[2]
		(ii)	The rate of this reaction is measured by recording the volume of gas produced every 30 seconds.
			Draw a labelled diagram to show the apparatus used to collect and measure the volume of gas produced. Your diagram must show how the apparatus is arranged.

[2]

(c) Fig. 10.1 is a graph to show the results when calcium carbonate and dilute hydrochloric acid react at room temperature.

The experiment is repeated with the dilute hydrochloric acid at a higher temperature. All other conditions remain the same.

Sketch on Fig. 10.1 the graph of the results for the experiment at a higher temperature of acid.



(d) The reaction between calcium carbonate and dilute hydrochloric acid is exothermic. Energy is transferred to the surroundings during this reaction.

[2]

Describe in terms of bond breaking and bond forming why a reaction is exothermic.	
[[3]
[Total: 1	0]

- 11 The isotope of sulfur $^{38}_{16}$ S decays to an element X by emitting a β -particle. Symbol X is **not** the chemical symbol of the element.
 - (a) (i) Complete the equation showing this process.

$$\frac{38}{16}S \rightarrow \dots X + \dots \beta$$
 [2]

(ii) Complete the sentence by using the Periodic Table on page 24 to identify element X.

Element X is[1]

(b) A scientist investigates the half-life of the isotope $^{38}_{16}$ S. Table 11.1 shows results from the scientist's notebook.

Table 11.1

time/h	reading on detector /s	corrected reading /s
0	42	40
0.5	37	35
1.0	32	30
1.5	29	27
2.0	26	24
2.5	23	21
3.0	21	19
3.5	18	16
4.0	16	14

(i) Deduce the background count.

		background count = / s [1]
(ii)	State one source of background radiation.	
		[11]

(iii) Use Table 11.1 to estimate the half-life of $^{38}_{16}$ S.

half-life =h [2]

[Total: 7]

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

The Periodic Table of Elements

	=	5 Z	helium 4	10	Ne	neon 20	18	Ar	argon 40	36	궃	krypton 84	54	×e	xenon 131	98	R	radon			
				6	ட	fluorine 19	17	Cl	chlorine 35.5	35	Ŗ	bromine 80	53	П	iodine 127	85	Αt	astatine -			
	>			8	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	<u>a</u>	tellurium 128	84	Ро	polonium –	116	^	livermorium –
	>			7	z	nitrogen 14	15	₾	phosphorus 31	33	As	arsenic 75	51	Sp	antimony 122	83	Ξ	bismuth 209			
	2			9	O	carbon 12	14	Si	silicon 28	32	Ge	germanium 73	50	Sn	tin 119	82	Pb	lead 207	114	Fl	flerovium
	=			2	В	boron 11	13	Αl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	<i>1</i> L	thallium 204			
										30	Zu	zinc 65	48	р О	cadmium 112	80	Hg	mercury 201	112	C	copernicium -
										29	Cn	copper 64	47	Ag	silver 108	62	Αn	gold 197	111	Rg	roentgenium -
Group										28	Z	nickel 59	46	Pd	palladium 106	78	瓧	platinum 195	110	Ds	darmstadtium -
Ğ				_						27	ဝိ	cobalt 59	45	몬	rhodium 103	77	'n	iridium 192	109	₹	meitnerium -
		- ⊐	hydrogen 1							26	Ьe	iron 56	44	Ru	ruthenium 101	9/	Os	osmium 190	108	Hs	hassium -
										25	Mn	manganese 55	43	ပ	technetium -	75	Re	rhenium 186	107	Bh	bohrium –
					pol	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
			Kev	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	g	niobium 93	73	Б	tantalum 181	105	ОР	dubnium –
					atc	nek				22	j=	titanium 48	40	Zr	zirconium 91	72	Ξ	hafnium 178	104	峜	rutherfordium -
										21	လွ	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89–103	actinoids	
	=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ഗ്	strontium 88	56	Ba	barium 137	88	Ra	radium
	_			8	=	lithium 7	17	Na	sodium 23	19	¥	potassium 39	37	Rb	rubidium 85	55	Cs	caesium 133	87	ᇁ	francium -

71 U	lutetium 175	103	۲	lawrencium	Ι
02 X	ytterbium 173	102	8	nobelium	Ι
69 L	thulium 169	101	Md	mendelevium	1
68 F	erbium 167	100	Fm	ferminm	Ι
²⁹	holmium 165	66	Es	einsteinium	Ι
99	dysprosium 163	86	ŭ	californium	I
65 Tb	terbium 159	26	Ř	berkelium	_
64 Gd	gadolinium 157	96	Cm	curium	I
63 Eu	europium 152	95	Am	americium	I
Sm	samarium 150	94	Pu	plutonium	Ι
61 Pm	promethium -	93	ď	neptunium	I
09 Z	neodymium 144	92	\supset	uranium	238
59 P	praseodymium	91	Ра	protactinium	231
58 G	cerium 140	06	H	thorium	232
22	lanthanum 139	68	Ac	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).