

Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

240475467

COMBINED SCIENCE

0653/42

Paper 4 Theory (Extended)

May/June 2022

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

This document has 20 pages.

DC (RW/JG) 302790/3 © UCLES 2022

[Turn over

1 (a) Fig. 1.1 shows a healthy human heart and a damaged human heart.




Fig. 1.1

- (i) Draw a label line and the letter **A** to show the position of the aorta on the healthy heart in Fig. 1.1. [1]
- (ii) The arrows on each heart show the direction of blood flow in the left side of the heart when the ventricles contract.

Identify one piece of evidence in Fig. 1.1 that shows the damaged heart has a faulty valve.
[1]
Coronary heart disease damages the heart. Diet is one risk factor of coronary heart disease.
State two other risk factors of coronary heart disease.
1
2

[2]

© UCLES 2022 0653/42/M/J/22

(iii)

(b) Fig. 1.2 is a graph showing the effect of physical activity on heart rate.




Fig. 1.2

(i) Calculate the percentage increase in heart rate between starting and stopping physical activity.Give your answer to the nearest whole number.

	% [3 _]
(ii)	Explain the change in heart rate between 30 and 40 minutes in Fig. 1.2.
	[3]

[Total: 10]

- 2 A student investigates a solid, a liquid and a gas.
 - (a) Three syringes contain 25 cm³ of either the solid, the liquid or the gas at room temperature and pressure, as shown in Fig. 2.1.

The end of each syringe is sealed.




Fig. 2.1

(i) The student pushes on the plunger of each syringe to increase the pressure.

The results are shown in Table 2.1.

Table 2.1

contents of syringe	volume at the start /cm ³	volume when pressure is increased / cm ³
solid	25	25
liquid	25	25
gas	25	21

Explain the results for each syringe when the pressure is increased.
Use ideas about particles in your answer.

	(ii)	ii) In a separate experiment, the student gently increases the temperature of the syringe that contains the gas, without pushing on the plunger.				
The volume of the gas changes from 25 cm ³ to 30 cm ³ .						
		Explain why the				
		Use ideas about	particles in your answer.			
					[2]	
(b)			solid X and a liquid Y separa C rise in temperature.	tely in two test-tubes and recor	ds the state	
	Tab	le 2.2 shows the re	esults.			
			Table 2.2			
		temperature /°C	state of X	state of Y		
		20	solid	liquid		
		25	solid	liquid		
		30	solid	liquid		
		35	solid	liquid		
		40	liquid	liquid		
		45	liquid	liquid		
	(i)	Use Table 2.2 to	estimate the melting point of	f X .		
			°C		[1]	
	(ii)	State one conclu	sion that can be made abou	t the boiling point of Y .		
					[1]	
(c)	The	solid X used in th	e experiment can burn.			
	Exp	olain why burning i	s a chemical change and me	elting is not a chemical change.		
					[11	

https://xtremepape.rs/

3 Fig. 3.1 shows the forces acting as a student rides forwards on a moving scooter.

The scooter has an electric motor.





Fig. 3.1

(a)	When the student is standing with both feet on the scooter, force ${\bf Q}$ is 340 N
	State the magnitude of force S .

Explain your answer.

force S =N	
explanation	
	[1]

- **(b)** The electric motor pushes the scooter forward with a constant force of 225 N for a distance of 0.30 m.
 - (i) Complete the boxes to show the useful energy transfers taking place.



(ii) Calculate the work done on the scooter by the electric motor.

(iii) The 225N force is applied for 1.2s.

Use your answer to (b)(ii) to calculate the useful power supplied to the scooter.

[Total: 7]

[2]

4 (a) Fig. 4.1 is a diagram of a wind-pollinated flower.




Fig. 4.1

Table 4.1 shows some of the labelled parts in Fig. 4.1 and their function.

Complete Table 4.1.

Table 4.1

letter	name of part	function
A		
D		
D		produce ovules
	stigma	

[3]

(b)	State one similarity and one difference between fertilisation in plants and fertilisation humans.	in
	similarity	
	difference	
		 [2]
(c)	After fertilisation in humans, a placenta develops inside the female uterus.	
	Describe how the placenta protects the fetus and allows it to grow.	
		[3]
	[Total:	8]

5 Fig. 5.1 shows a key made from the alloy brass.




Fig. 5.1

(a)	Suggest two reasons why brass is more suitable for making keys than pure copper.	
	1	
	2	
		[2]

(b) Brass contains copper atoms and zinc atoms, as shown in Fig. 5.2.




Fig. 5.2

Table 5.1 shows some information about brass.

Table 5.1

	percentage of element in brass	relative size of atom
copper	56	1.0
zinc	37	1.1

Fig. 5.2 is **not** an accurate representation of brass because brass contains more than two elements.

(i)	Describe how the information in Table 5.1 shows that there are more than two element in brass.
	[1

(ii)	State two other reasons why Fig. 5.2 is not an accurate representation of the atoms i brass.
		Use Table 5.1 to help you.
		1
		2
		[2
(c) /	A st	udent investigates the reactivity of copper and zinc.
		student places a zinc rod into a solution containing aqueous copper ions and leaves it for inutes, as shown in Fig. 5.3.
		appearance at the start appearance after 5 minutes
	gre	rod has orange coating
С		paler blue per(II) ions
		Fig. 5.3
((i)	State why the zinc rod has an orange coating after 5 minutes.
		[1
(ii)	State why the colour of the solution changes.
		[1
(i	ii)	In another experiment, a copper rod is placed into an aqueous solution of zinc ions.
		Describe the appearance of the rod and the solution after 5 minutes.
		Explain your answer.

- 6 Ultraviolet radiation and microwaves are part of the electromagnetic spectrum.
 - (a) Fig. 6.1 shows an incomplete electromagnetic spectrum.

On Fig. 6.1, write ultraviolet and microwaves in the correct places.

increasing frequency						
gamma radiation			visible light			radio waves

Fig. 6.1		

		[2]
(b)	State one danger of ultraviolet radiation.	
		[1]
(c)	State one use of microwaves.	
		[1]

(d) Complete the sentences about ultraviolet radiation and microwaves.

Circle the correct word or phrase to complete each sentence.

Ultraviolet radiation and microwaves are **audible / longitudinal / transverse** waves.

The speed of microwaves in a vacuum is **equal to** / **faster than** / **slower than** the speed of ultraviolet radiation in a vacuum.

[2]

(e) (i) The speed of ultraviolet radiation in a vacuum is 3.0×10^8 m/s.

An ultraviolet lamp emits ultraviolet radiation of wavelength $3.5 \times 10^{-7} \, \text{m}$.

Calculate the frequency of ultraviolet radiation at this wavelength.

Give the unit of your answer.

(ii)	Three identical ultraviolet lamps are connected in parallel to a 230 V electricity supply.
	Each lamp uses a power of 150 W.
	Calculate the total current from the electricity supply.

current =		Α	[3]
-----------	--	---	-----

[Total: 12]

7 (a) Fig. 7.1 shows a food chain.




Fig. 7.1

		J	
	(i)	Identify the primary consumer in Fig. 7.1.	
			[1]
	(ii)	Explain why food chains usually have fewer trophic levels than the food chain in Fig. 7	'.1.
			[2]
			[~]
(b)	Gra	ss is a type of plant.	
	(i)	The leaves of a plant have different types of cells with different functions.	
		State the main function of the two types of mesophyll cells.	
		palisade mesophyll cells	
		spongy mesophyll cells	
			[2]
	(ii)	Plant growth can be affected by ion deficiencies.	
		Explain the effects of magnesium ion deficiency on plant growth.	
			$\Gamma \cap I$

(c)	In humans, food is digested in the stomach of the alimentary canal. Hydrochloric acid is found in the stomach.				
	State two functions of hydrochloric acid in the stomach.				
	1				
	2				
		[2]			

[Total: 9]

8 Energy level diagrams for two reactions are shown in Fig. 8.1. The diagrams are drawn to the same scale.




Fig. 8.1

(a)	(i)	Compare the activation energies for reaction 1 and reaction 2.	
		Explain your answer.	
		comparison	
		explanation	
			 [1]
	(ii)	The temperature of reaction 2 is increased.	
		Use ideas about activation energy to explain why this increases the rate of reaction 2.	
			[1]
(b)	One	e of the reactions represents the combustion of propane.	
	lder	ntify which reaction, 1 or 2, represents the combustion of propane.	
	Give	e a reason for your answer.	
	read	ction	
	reas	son	
			 [1]

(c) Arrows A and B on Fig. 8.2 represent changes occurring during reaction 1.




Fig. 8.2

Put ticks (\checkmark) in the boxes to show what these changes represent.

	bonds are being broken	bonds are being formed	energy is being taken in	energy is being given out
arrow A				
arrow B				

[2]

(d) Equations for the combustion of propane are shown in Fig. 8.3.

Fig. 8.3

State the number of electrons that are shared between one oxygen atom and one carbon atom in a molecule of carbon dioxide.

1	11
 1.	

(ii) State **two** ways in which the structure of propane shows that it is an alkane.

1	
2	
	[2]

[Total: 8]

9 (a) Fig. 9.1 shows a liquid-in-glass thermometer without a scale.



Fig. 9.1

The thermometer measures temperatures between -10 °C and +110 °C.

Table 9.1 gives some information about four liquids, **A**, **B**, **C** and **D**.

Table 9.1

liquid	melting point /°C	boiling point /°C
Α	-89	+117
В	-117	+79
С	-39	+367
D	+17	+118

State the letters of the liquids in Table 9.1 that can be used in this thermometer.

_______[1]

(b) Fig. 9.2 shows a circuit used to heat a beaker of water with an electric heater.




Fig. 9.2

The circuit contains a variable resistor.

	Draw the circuit symbol for the variable resistor.
	[1]
(ii)	The variable resistor is adjusted so that the current is 5.0A.
	Calculate the energy in kilojoules supplied by the 12V battery when the circuit is switched on for 5 minutes.
	energy =kJ [3]
(iii)	A voltmeter connected across the variable resistor shows a reading of 4.0 V.
	Calculate the potential difference across the electric heater.
	potential difference =V [1]
(iv)	Some electrical energy supplied by the battery is not transferred to useful thermal energy in the water.
	Suggest how electrical energy from the battery is lost in the circuit.
	[2]
	[Total: 8]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

ts
Ī
Je
e
Ш
o
<u>a</u>
ap
Ë
<u>ပ</u>
Q
<u>ŏ</u>
ē
₫.
he
\vdash

	=	2 He	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	R	radon			
	=>			6	ш	fluorine 19	17	Cl	chlorine 35.5	35	Ä	bromine 80	53	Н	iodine 127	85	At	astatine			
	5			80	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	Б	tellurium 128	84	Ъ	polonium -	116	_	livermorium -
	>			7	z	nitrogen 14	15	۵	phosphorus 31	33	As	arsenic 75	51	Sp	antimony 122	83	<u>.</u>	bismuth 209			
	≥			9	ပ	carbon 12	14	S	silicon 28	32	Ge	germanium 73	92	Sn	tin 119	82	Ъ	lead 207	114	Εl	flerovium —
	=			2	В	boron 11	13	Αl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	11	thallium 204			
										30	Zu	zinc 65	48	g	cadmium 112	80	Нg	mercury 201	112	ပ်	copemicium
										29	Cn	copper 64	47	Ag	silver 108	79	Αn	gold 197	111	Rg	roentgenium -
Group										28	Z	nickel 59	46	Pq	palladium 106	78	础	platinum 195	110	Ds	darmstadtium -
) Dig										27	ပိ	cobalt 59	45	格	rhodium 103	77	٦	iridium 192	109	M	meitnerium -
		- I	hydrogen 1							26	Ьe	iron 56	44	Ru	ruthenium 101	92	Os	osmium 190	108	Hs	hassium -
										25	Mn	manganese 55	43	ည	technetium -	75	Re	rhenium 186	107	Bh	bohrium –
					pol	ass				24	ပ်	chromium 52	42	Мо	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
			Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	qN	niobium 93	73	ц	tantalum 181	105	Οþ	dubnium —
					ato	rels				22	F	titanium 48	40	Zr	zirconium 91	72	Ξ	hafnium 178	104	Ŗ	rutherfordium —
										21	Sc	scandium 45	39	>	yttrium 89	57–71	lanthanoids		89–103	actinoids	
	=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	Š	strontium 88	56	Ba	barium 137	88	Ra	radium -
	_			က	:=	lithium 7	+	Na	sodium 23	19	¥	potassium 39	37	&	rubidium 85	55	S	caesium 133	87	Ϋ́	francium -

71 Lu	lutetium 175	103	۲	lawrencium
70 Yb	ytterbium 173	102	Š	nobelium —
mL Tm	thulium 169	101	Md	mendelevium —
68 Er	erbium 167	100	Fm	fermium —
67 H0	holmium 165	66	Es	einsteinium —
% Dy	dysprosium 163	86	Ç	califomium —
65 Tb	terbium 159	97	Ř	berkelium —
64 G d	gadolinium 157	96	Cm	curium —
63 Eu	europium 152	92	Am	americium -
62 Sm	samarium 150	94	Pn	plutonium —
61 Pm	promethium -	93	Ν	neptunium —
9 P N	neodymium 144	92	\supset	uranium 238
59 Pr	praseodymium 141	91	Ра	protactinium 231
58 Ce	cerium 140	06	드	thorium 232
57 La	lanthanum 139	88	Ac	actinium —

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).