

Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

202719202

COMBINED SCIENCE

0653/63

Paper 6 Alternative to Practical

October/November 2022

1 hour

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].

This document has 16 pages. Any blank pages are indicated.

1 (a) Fig. 1.1 shows a photograph of a section of a star fruit (carambola) containing three seeds.




Fig. 1.1

(i)	The distance between points P and Q on Fig. 1.1 represents the width of the star fruit.
	Measure the width PQ on Fig. 1.1.

(ii) The actual width of the star fruit is 80 mm.

Calculate the magnification of the photograph in Fig. 1.1.

Use the equation shown.

magnification =
$$\frac{\text{width of } \mathbf{PQ} \text{ in Fig. 1.1}}{\text{actual width of star fruit}}$$

(iii)	In the box, make a large, detailed pencil drawing of the star fruit shown in Fig. 1.1

[3]

(iv) On your drawing in (a)(iii), add a label line and the letter S to show one of the seeds. [1]

(b) A student investigates the nutrient content of the star fruit.

Procedure

The student:

- puts a small sample of the star fruit into each of four test-tubes
- adds a different test solution to each test-tube
- shakes the test-tubes and observes the results.

Table 1.1 shows the test solutions used and the colour after testing.

Table 1.1

test-tube	test solution used	colour of test solution before testing	colour of test solution after testing	conclusion
1	iodine solution		blue-black	
2	Benedict's solution		blue	no reducing sugar present
3	biuret solution		blue	
4	universal indicator	green	red	

(i)	Complete the third column of Table 1.1 by adding the colour of the test solution before testing.	ons [3]
(ii)	Complete the last column of Table 1.1 by stating a conclusion for each test.	[3]
iii)	State why the conclusion for the Benedict's test may not be correct.	
		[1]

[Total: 13]

2 A student investigates the effect of heat on solid sodium hydrogencarbonate.

Procedure

The student:

- adds 1.0 g of solid sodium hydrogencarbonate to a test-tube
- · heats the sodium hydrogencarbonate using the blue flame of a Bunsen burner for five minutes
- allows the test-tube to cool down
- measures the mass of the solid remaining in the test-tube.

The student repeats the procedure using different masses of sodium hydrogencarbonate.


The results are shown in Table 2.1

Table 2.1

mass of sodium hydrogencarbonate heated /g	mass of solid remaining in the test-tube /g
1.0	0.8
2.0	2.7
3.0	2.5
4.0	3.4
5.0	4.2
6.0	
7.0	

(a) Fig. 2.1 shows the balance readings for 6.0 g and 7.0 g of sodium hydrogencarbonate heated.








7.0 g of sodium hydrogencarbonate

Fig. 2.1

Record in Table 2.1 these readings to **one** decimal place.

[2]

(b) (i) On the grid, plot a graph of mass of solid remaining in the test-tube (vertical axis) against mass of sodium hydrogencarbonate heated.



(ii)	Circle the anomalous point on the graph.	[1]
(iii)	Draw the best-fit straight line.	[1]
(iv)	Describe the relationship between the mass of solid remaining in the test-tube and mass of sodium hydrogencarbonate heated.	l the
(v)	Use your graph to estimate the mass of solid remaining in the test-tube when 4.7	
(*)	solid sodium hydrogencarbonate is heated.	y o
	mass of solid remaining in the test-tube =	ı [1]

[3]

(C)		n a gas.
	(i)	Explain how the results in Table 2.1 support this conclusion.
		[1]
	(ii)	The student thinks that the gas given off is carbon dioxide.
		Describe the test for carbon dioxide and state the observation for the positive result.
		test
		observation[1]
(d)	_	gest why the student heats the test-tube with the blue flame of a Bunsen burner instead ne yellow flame.
		[1]
(e)		student thinks there is still some unreacted sodium hydrogencarbonate left in the tube after heating for five minutes.
	_	gest what the student does to test if there is any unreacted sodium hydrogencarbonate in the test-tube.
		[1]
		[Total: 13]

A st	udent determines the thermal energy lost by a beaker of water when an ice cube melts in the cuber.	he					
(a)	The student measures 100 cm ³ of water and pours it into a beaker.						
	Suggest a suitable piece of apparatus to measure 100 cm ³ of water.	[1]					
(b)	The student measures the initial temperature T_i of the water.						
	Fig. 3.1 shows the thermometer reading.						
	Fig. 3.1 Record the initial temperature T_i of the water to the nearest 0.5 °C.						
(0)	T _i =°C	[1]					
(6)	Procedure						
	The student:						
	 adds an ice cube to the beaker of water waits for the ice cube to melt stirs the water measures the final temperature T_f of the water. 						
	Suggest why stirring the water improves the accuracy of the measurement of $T_{\rm f}$.						
		[1]					

(d)	The final temperature $T_f = 10.5$ °C.
	Calculate the change in temperature ΔT of the water in the beaker.
	Use your answer in (b) and the equation shown.
	$\Delta T = T_{\rm i} - T_{\rm f}$
	ΔT =°C [1]
(e)	Calculate the thermal energy \boldsymbol{E} lost by the water in the beaker as the ice cube melts.
	Use the equation shown.
	$E = 420 \times \Delta T$
	Give your answer to two significant figures.
	<i>E</i> = J [2]
	⊏ J [2]
(f)	The thermal energy lost by the water is less than the thermal energy needed to melt the ice cube.
	Suggest where the extra thermal energy comes from.
	[1]
	[Total: 7]

4 A student wants to determine whether the resistance of a wire is related to the metal used to make the wire.

The student has wires made of three different metals: constantan, nichrome and tungsten.

The resistance *R* of a wire is calculated using the equation:

$$R = \frac{V}{I}$$

where V is the potential difference (p.d.) across the wire and I is the current in the wire.

Plan an investigation to determine whether the resistance of a wire is related to the metal used to make the wire.

You are provided with:

- constantan wires of different lengths and thicknesses
- nichrome wires of different lengths and thicknesses
- tungsten wires of different lengths and thicknesses.

You may use any common laboratory apparatus in your plan.

Include in your plan:

- the apparatus you will use, you may include a circuit diagram
- a brief description of the method, including the measurements you will make
- which variables you will keep constant
- a results table to record your measurements (you are not required to enter any readings in the table)
- how you will process your results to draw a conclusion.

			[7]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.