

Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

4077882846

CO-ORDINATED SCIENCES

0654/43

Paper 4 Theory (Extended)

May/June 2022

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 120.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

This document has 36 pages. Any blank pages are indicated.

DC (CE) 303339/2 © UCLES 2022

[Turn over

1 (a) Fig. 1.1 is a diagram of the female reproductive system.




Fig. 1.1

	Identify the letter from Fig. 1.1 that repr	esents:	
	where female gametes are released		
	where fertilisation occurs		
	where implantation occurs		
	where meiosis occurs.		[4]
(b)	Female gametes in humans are called	egg cells.	
	State the name of the male gamete in h	numans.	
			[1

(c) Table 1.1 compares some features of male and female gametes.

Complete Table 1.1.

Table 1.1

	male gamete	female gamete
relative size		
number released at one time		usually one
motility		

[3]

(d) Complete the sentences to describe the adaptive features of egg cells.

One of the adaptive features of egg cells is that it has stores.

The egg cell also has a coating that changes after fertilisation. [2]

[Total: 10]

2 Petroleum is a fossil fuel.

It can be separated into useful fractions by fractional distillation.

Fig. 2.1 shows a diagram of a fractionating column.




Fig. 2.1

(a)	(i)	Explain why it is possible to separate the substances in petroleum by fractional distillation	on.
			 [1]
	(ii)	On Fig. 2.1, write the letter X in the coolest part of the fractionating column.	[1]
(b)	Tab	le 2.1 shows the uses of some of the fractions.	
	Con	nplete Table 2.1.	

Table 2.1

fraction	use
refinery gas	bottled gas for heating
gasoline	
naphtha	feedstock for making chemicals
diesel oil	
bitumen	

[3]

(c) Refinery gas contains propane, C₃H₈.

	Draw a diagram to show the structure of propane.	
		[2]
(d)	Refinery gas also contains butane, C ₄ H ₁₀ .	

Butane burns in oxygen to make carbon dioxide and water.

Construct the balanced symbol equation for this reaction.

[2]

(e) Burning butane is a chemical change.

Describe the difference between a chemical change and a physical change.

[1]

[Total: 10]

3 (a) Fig. 3.1 shows a piece of graphite with an irregular shape.




Fig. 3.1

	(i)	Describe a method to determine the volume of the piece of graphite.
		[2]
	(ii)	The piece of graphite has a mass of 33 g and a volume of 15 cm ³ .
		Calculate the density of the piece of graphite.
		density = g/cm ³ [2]
(b)	Gra	phite can be used as a lubricant in machines with moving parts such as an electric drill.
	(i)	Describe, in terms of forces and energy transfers, how lubricants increase the efficiency of a machine.
		[3]

(ii)	An electric drill transfers 1200 J of electrical energy to 900 J of useful kinetic energy.
	Calculate the efficiency of the electric drill.
	
	efficiency = % [2]
(iii)	The electric motor in the drill has a current of 25A when using an 18V battery.
	Calculate the power output of the motor.
	power = W [2]
	[Total: 11]

4 A student investigates the effect of temperature on the rate of photosynthesis.

Fig. 4.1 shows the apparatus they use.




Fig. 4.1

The student counts the number of oxygen bubbles produced in one minute.

He repeats this investigation, changing the temperature of the water each time.

The number of oxygen bubbles produced per minute is equivalent to the rate of photosynthesis.

Table 4.1 shows the results.

Table 4.1

temperature/°C	number of bubbles produced per minute
0	0
5	4
10	8
15	13
20	16
25	18
30	19
35	8
40	0

(a)	State the temperature from	Table 4.1	where the	rate of	photosynthesis	is the	highest
-----	----------------------------	-----------	-----------	---------	----------------	--------	---------

00	[4]
	- 1 1 1

(b)	Pho	otosynthesis is an enzyme-controlled reaction.	
	(i)	Explain the results in Table 4.1 between 5 °C and 15 °C.	
			[3]
	(ii)	State the temperature from Table 4.1 when all the enzymes involved in photo are completely denatured.	osynthesis
			°C [1]
(c)	The	carbohydrate glucose is also a product of photosynthesis.	
	Glu	cose is converted to different substances for transport and storage in a plant.	
	(i)	Describe how carbohydrates are transported in a plant.	
			[3]
	(ii)	State the larger molecule made from glucose that is used for storage in a plan	t.
			[1]
(d)	Chlo	orophyll is also necessary for photosynthesis.	
	Stat	te the energy transfer that chlorophyll is responsible for.	
		energy → energy	[2]
			[Total: 11]

- 5 This question is about electrolysis.
 - (a) The list shows the particles found in aqueous copper(II) sulfate.

 Cu^{2+} H^{+} $H_{2}O$ SO_{4}^{2-} OH^{-}

State the formula of **one** particle attracted to the **cathode** during electrolysis.

Choose from the list.

.....

(b) Aqueous copper(II) sulfate conducts electricity.

Explain why.

[1]

(c) Fig. 5.1 shows the apparatus used for the electrolysis of aqueous copper(II) sulfate.




Fig. 5.1

0654/43/M/J/22

(i) State the name given to the **positive** electrode.

......[1]

© UCLES 2022

	(ii)	The purification (refining) of copper uses electrolysis.
		Describe how impure copper is purified by electrolysis.
		Include ionic half-equations in your answer.
		[4]
(d)	Loo	k at this ionic half-equation.
		$Al^{3+} + 3e^{-} \rightarrow Al$
	Stat	te if this reaction is an example of oxidation or reduction.
	Ехр	lain your answer.
		[1]
		[Total: 9]

6 Fig. 6.1 shows a cheetah.

Cheetahs are the fastest land animal and have a top speed of 30 m/s.




Fig. 6.1

(a)	State the difference between speed and velocity.
	r.a

(b) Fig. 6.2 shows a speed–time graph for a cheetah's journey.




Fig. 6.2

Describe the motion of the cheetan shown in Fig. 6.2.	
Г	2

(c)	The	mass of the cheetah is 42 kg.
	Cal	culate the kinetic energy of the cheetah when it is running at its maximum speed of 30 m/s.
		kinetic energy = J [2]
		Kirietic eriergy –
(d)	A ch	neetah drinks water from a puddle.
	Ove	er time, the water in the puddle evaporates.
	Eva	poration and boiling both turn liquid water into a gas.
	(i)	State one difference between evaporation and boiling.
		[1]
	(ii)	State two ways to increase the rate of evaporation from the puddle.
		1
		2
		[2]
		[Total: 0]

[Total: 9]

7 (a) The effect of an injection of adrenaline on pulse rate is recorded.

The adrenaline is injected at 1 minute.

Fig. 7.1 shows a graph of the results.




Fig. 7.1

	((i)	Identify	/ in	Fig.	7.	1 the	pulse	rate	before	the	adrenaline	in (jection
--	---	-----	----------	------	------	----	-------	-------	------	--------	-----	------------	------	---------

	beats per minute [1
(ii)	Describe the immediate effect of the adrenaline injection on pulse rate shown in Fig. 7.1
	Use data from the graph to support your answer.
	[2
(i)	State one effect adrenaline has on the eye.
	[1
(ii)	Name the nerve that carries impulses from the eye to the brain

© UCLES 2022 0654/43/M/J/22

(b)

(c)	A ho	ormone decreases blood glucose concentration by causing the glucose to be stored.
	(i)	State the name of the hormone that decreases blood glucose concentration.
		[1]
	(ii)	State the name of the organ that produces this hormone.
		[1]
	(iii)	State the name of the organ that stores the excess glucose.
		[1]
(d)	Des	cribe two ways the actions of the hormonal system are different from the nervous system.
	1	
	2	
		[2]
		[Total: 10]

8 Fig. 8.1 shows the arrangement of ions in magnesium metal at 25°C.




Fig. 8.1

(a)	Describe the changes in the arrangement and movement of magnesium ions when magnesium melts.
	changes in arrangement of magnesium ions
	changes in movement of magnesium ions
	[2]
(b)	Magnesium melts at 650 °C and boils at 1090 °C.
	In the box, draw the arrangement of ions in magnesium at 1800 °C.

(c) Magnesium reacts with dilute hydrochloric acid. Hydrogen gas is made in the reaction.

A student investigates this reaction. Fig. 8.2 shows the apparatus he uses.




Fig. 8.2

Every 10 seconds, the student measures the total volume of hydrogen gas made.

Fig. 8.3 shows the graph the student plots of his results.




Fig. 8.3 0654/43/M/J/22

	(i)	State the volume of gas collected after 40 seconds.
		volume of gas = cm ³ [
	(ii)	The reaction is fastest during the first 10 seconds.
		Explain why.
		[
	(iii)	The student repeats the experiment.
		He uses the same volume of hydrochloric acid and the same mass of magnesium.
		This time he increases the temperature of the hydrochloric acid.
		All of the magnesium reacts with the acid.
		On Fig. 8.3, sketch the shape of the graph you would expect this time.
(d)		e rate of the reaction can be increased by increasing the concentration of the dilut rochloric acid.
	Exp	olain why. Use ideas about collisions between particles.
		[2
(e)	Mag acid	gnesium chloride is also made in the reaction between magnesium and dilute hydrochlorid.
	Ма	gnesium chloride contains the ions ${ m Mg}^{2+}$ and ${ m C}\it{l}^-$.
	Det	ermine the formula of magnesium chloride.
		[
		[Total: 10

9 A student investigates the effect of changing light levels on the resistance of a light-dependent resistor (LDR).

The student shines a torch (flashlight) on to the LDR.

She then places glass slides between the LDR and the torch (flashlight) to reduce the light intensity (amount of light) reaching the LDR.

Fig. 9.1 shows the equipment she uses.




Fig. 9.1

The student places more glass slides between the torch (flashlight) and the LDR and measures the resistance, in kilo-ohms ($k\Omega$), using a resistance meter.

(a) Fig. 9.2 shows a graph of the student's results.




Fig. 9.2

(i)	Use Fig. intensity.	9.2	to	describe	how	the	resistance	of	the	LDR	varies	with	changing	ligh
														. [2
(ii)	The resis	tance	m د	eter nrovi	ides a	a not	ential differe	2nc	e (n	d) of	14 V a	cross	the I DR	

Calculate the charge flowing through the LDR in 1 minute when 3 glass slides are used.

charge = C [4]

(b)	The	lamp emits visible light at a frequency of 5.0×10^{14} Hz.
	(i)	State the meaning of the word frequency.
		[1]
	(ii)	Calculate the wavelength of this visible light.
		wavelength = m [3]
	(iii)	State one form of electromagnetic radiation that has a frequency higher than visible light.
		[1]
		[Total: 11]

10 (a) The blood group of some patients in hospital is recorded.

Fig. 10.1 shows a bar chart of the results.




Fig. 10.1

(i)	One of the blood groups in Fig. 10.1 is not labelled.
	State this blood group.
	[1]
(ii)	Identify the most common blood group in Fig. 10.1.
	[1]
(iii)	Describe evidence from Fig. 10.1 that shows that this characteristic is an example of discontinuous variation.
	[2]
(iv)	State the cause of the variation seen in Fig. 10.1.

(b) Adaptations in populations can be inherited through natural selection or selective breeding.
 Table 10.1 compares some features of natural selection and selective breeding.
 Complete Table 10.1 by placing ticks (✓) in the boxes to show the correct features.

Table 10.1

	involves passing on of alleles to offspring	is used to improve domesticated animals	occurs over many generations	keeps the features best suited to the environment
natural selection				
selective breeding				

[2]

Sulf	furic	acid is made by the Contact process.									
Sulf	Sulfur, air and water are raw materials used to make sulfuric acid.										
Loo	k at	the equations for the first two stages in the Contact process.									
sta	ge 1	+ sulfur dioxide									
sta	ge 2	sulfur dioxide + oxygen ⇌ sulfur trioxide									
(a)	Cor	nplete the word equation for stage 1 of the Contact process. [2]									
(b)	The	conditions used for stage 2 are:									
	•	450 °C atmospheric pressure a catalyst.									
	(i)	State the name of the catalyst used.									
		[1]									
	(ii)	Explain why a catalyst and a temperature of 450 °C are used in stage 2 of the Contact process.									
		Use ideas about:									
		 the percentage of sulfur trioxide made the rate of reaction. 									
		catalyst									
		percentage of sulfur trioxide made									
		rate of reaction									
		450°C									
		percentage of sulfur trioxide made									
		rate of reaction									
		[4]									

© UCLES 2022 0654/43/M/J/22

11

(c) The reaction in stage 2 of the Contact process is exothermic.

Fig. 11.1 shows the energy level diagram for the reaction.

Complete the labels on Fig. 11.1.

Choose the labels from the list.

energy given out energy taken in products reactants

reactants have less energy




Fig. 11.1

[2]

(d)	In an experiment, 200 g of sulfur trioxide, SO ₃ , is made.
	Calculate the volume occupied by 200 g of sulfur trioxide gas.
	The relative molecular mass, $M_{\rm r}$, of sulfur trioxide is 80.
	The volume of one mole of any gas is 24 dm ³ at room temperature and pressure (r.t.p.).
	Show your working.

 $volume = \dots dm^3 [2]$

[Total: 11]

12 Fig. 12.1 shows a forklift truck lifting a crate.




Fig. 12.1

(a) The forklift truck does 2750 J of work on the crate when the crate is lifted through a height of 2.2 m.

The gravitational field strength, g, is 10 N/kg.

Calculate the mass of the crate.

mass = kg [2]

(b) Fig. 12.2 shows the same forklift truck after it has lowered the crate.




Fig. 12.2

Explain why the forklift truck is more stable after it has lowered the crate.

Use ideas about centre of mass in your answer.

.....[1]

(c) The forklift truck uses an electric motor to lift the crate.

Fig. 12.3 shows a simple d.c. motor.




Fig. 12.3

		G
	(i)	A current flows through the coil.
		Draw arrows on Fig. 12.3 to show the direction of the force acting on points X and Z of the coil.
	(ii)	State why point Y does not experience a force.
		[
(d)	Α β-	particle passes between the poles of a permanent magnet.
	(i)	Suggest why a β -particle is deflected when moving through a magnetic field.
		[2
	(ii)	State and explain how the deflection direction of an α -particle would differ from that the β -particle.

[Total: 9]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

ts
Ĭ
ခ
ב
<u>•</u>
Ш
0
<u>o</u>
a
\vdash
ပ
믕
ŏ
÷.
ᇹ
~
_
<u>@</u>
ᆚ
_

	=	2 He	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton 84	25	Xe	xenon 131	98	Ru	radon			
	=			6	ш	fluorine 19	17	Cl	chlorine 35.5	35	Ŗ	bromine 80	53	Н	iodine 127	82	Ą	astatine -			
	5			80	0	oxygen 16	16	S	sulfur 32	34	Se	selenium 79	52	<u>P</u>	tellurium 128	84	Ъо	polonium –	116		livermorium —
	>			7	z	nitrogen 14	15	۵	phosphorus 31	33	As	arsenic 75	51	Sp	antimony 122	83	Ξ	bismuth 209			
	≥			9	ပ	carbon 12	14	S	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Ъ	lead 207	114	F1	flerovium
	≡			5	М	boron 11	13	Αl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	84	lΤ	thallium 204			
										30	Zu	zinc 65	48	ဗ	cadmium 112	80	Ą	mercury 201	112	S	copernicium
										59	Cn	copper 64	47	Ag	silver 108	62	Αn	gold 197	111	Rg	roentgenium -
Group										28	z	nickel 59	46	Pd	palladium 106	78	చ	platinum 195	110	Ds	darmstadtium -
Gro										27	රි	cobalt 59	45	牊	rhodium 103	77	Ir	iridium 192	109	M	meitnerium -
		- エ	hydrogen 1							26	Ьe	iron 56	44	Ru	ruthenium 101	9/	Os	osmium 190	108	Hs	hassium –
										25	Mn	manganese 55	43	ပ	technetium -	75	Re	rhenium 186	107	Bh	bohrium –
				_	pol	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
			Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	qN	niobium 93	73	д	tantalum 181	105	Db	dubnium –
					atc	rek				22	j	titanium 48	40	Zr	zirconium 91	72	Ξ	hafnium 178	104	Ŗ	rutherfordium -
										21	Sc	scandium 45	39	>	yttrium 89	57–71	lanthanoids		89–103	actinoids	
	=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	Š	strontium 88	56	Ba	barium 137	88	Ra	radium -
	_			က	=	lithium 7	1	Na	sodium 23	19	¥	potassium 39	37	ВВ	rubidium 85	55	S	caesium 133	87	ъ́	francium -

71	Γn	lutetium	175	103	۲	lawrencium	1
70	Υp	ytterbium	173	102	8	nobelium	ı
69	TB	thulium	169	101	Md	mendelevium	ı
89	Щ	erbinm	167	100	Fm	ferminm	ı
29	웃	holmium	165	66	Es	einsteinium	ı
99	ò	dysprosium	163	86	ŭ	californium	ı
65	Tp	terbium	159	97	BK	berkelium	1
64	В	gadolinium	157	96	Cm	curium	ı
63	En	europium	152	92	Am	americium	1
62	Sm	samarium	150	94	Pu	plutonium	ı
61	Pm	promethium	1	93	ď	neptunium	ı
	PΝ	ne					
69	Ā	praseodymium	141	91	Ра	protactinium	231
58	Ce	cerium	140	06	드	thorium	232
22	Гa	lanthanum	139	68	Ac	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is $24\,\mathrm{dm}^3$ at room temperature and pressure (r.t.p.).