

# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Ordinary Level

CANDIDATE NAME										
CENTRE NUMBER						CAN NUM				_

027921705

CHEMISTRY 5070/02

Paper 2 Theory

1 hour 30 minutes

May/June 2008

Candidates answer on the Question Paper.

Additional Materials: Answer Booklet/Paper

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do **not** use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

# **Section A**

Answer all questions.

Write your answers in the spaces provided in the Question Paper.

#### Section B

Answer any **three** questions.

Write your answers on any lined pages and/or separate answer paper.

A copy of the Periodic Table is printed on page 16.

The number of marks is given in brackets [ ] at the end of each question or part question.

At the end of the examination, fasten all your work securely together.

For Examiner's Use				
Section A				
В7				
В8				
В9				
B10				
Total				

This document consists of **16** printed pages.



# **Section A**

For Examiner's Use

[Total: 5]

Answer all the questions in this section in the spaces provided.

The total mark for this section is 45.

**A1** Choose from the following gases to answer the questions below.

ammonia argon carbon monoxide chlorine

hydrogen nitrogen

nitrogen dioxide

oxygen

Each gas can be used once, more than once or not at all.

Name a gas which

(a)	is made during the incomplete combustion of octane,
	[1]
(b)	dissolves in water to make an alkaline solution,
	[1]
(c)	is monatomic,
	[1]
(d)	is a reducing agent in a Blast Furnace,
	[1]
(e)	is used in the Contact process.
	[1]

© UCLES 2008 5070/02/M/J/08

<b>A2</b>	Iron(II) sulphate, FeSO <sub>4</sub> , is easily oxidised to iron(III) sulphate.										
	(a)	Cald	culate the percentage by mass of iron in iron(II) sulphate.								
			% [2]								
	(b)		ample of iron(II) sulphate is dissolved in water. Describe a test to show the presence ulphate ions in this solution.								
		rea	gents								
		obs	ervation[2]								
	(c)	In the presence of aqueous hydrogen ions and dissolved oxygen, aqueous iron(II) ions are oxidised to form iron(III) ions and water. Write an ionic equation for this reaction.									
		[2]									
	(d)	Aqu	leous iron(II) ions can also be oxidised by reaction with acidified potassium bromate(VI), $K_2Cr_2O_7$ . At the same time aqueous dichromate(VI) ions are reduced.								
		(i) Describe the colour change of the chromium-containing species during the reaction.									
			[1]								
		(ii)	Describe the colour change of the iron-containing species during the reaction.								
			[1]								

(e)	An i	mpure sample of iron(II) sulphate was analysed by titration.					
	The sample was dissolved in 25.0 cm <sup>3</sup> of dilute sulphuric acid and then titrate 0.0400 mol/dm <sup>3</sup> potassium dichromate(VI) solution.						
	19.0	cm <sup>3</sup> of potassium dichromate(VI) solution was required to reach the end-point.					
	(i)	Calculate the number of moles of potassium dichromate(VI) used in the titration.					
		moles [1]					
	(ii)	One mole of potassium dichromate(VI) reacts with six moles of iron(II) ions. Calculate the mass, in grams, of iron(II) ions in the sample analysed.					
		mass of iron(II) ions g [2]					
		[Total: 11]					

© UCLES 2008 5070/02/M/J/08

A3 A student found a copy of a Periodic Table published in the year 1930. Several elements we missing from this table because they had not yet been discovered. One of these elements was technetium, Tc.													
	One	ne isotope of technetium has the symbol $^{98}_{43}$ Tc.											
	(a)	(a) Complete the table below to show the number of subatomic particles in one atom of this isotope.											
			number of protons										
			number of electrons										
			number of neutrons										
					[2]								
	(b)	Suggest the sy	mbol of another isotope	e of technetium.									
					[1]								
	(c)	Explain, in terms of subatomic particles and their charge, why an atom of $^{98}_{43}{\rm Tc}$ is electrically neutral.											
	[												
	(d)	From its position	on in the modern Period	ic Table predict two prop	erties of technetium.								
		1											
		2			[2]								
					[Total: 7]								

© UCLES 2008 5070/02/M/J/08 **[Turn over** 

(a)	One, C <sub>2</sub> H <sub>6</sub> , and ethene, C <sub>2</sub> H <sub>4</sub> , are both gaseous hydrocarbons.							
aj	Describe how aqueous bromine can be used to distinguish between a sample of ethane and a sample of ethene.							
	[2]							
(b)	Draw a 'dot-and-cross' diagram for ethane. You only need to draw the outer electrons of the carbon atoms.							
	[2]							
(c)	Ethane reacts with chlorine in the presence of ultra-violet light.  Suggest a structure for a product of this reaction.							
	[1]							
(d)	Write both the name and the molecular formula of an alkene molecule containing four carbon atoms.							
	name							

© UCLES 2008 5070/02/M/J/08 A5 One of the largest uses of phosphorus is in the making of safety matches. A safety match ignites when it is rubbed against the striking surface of a match box. The match head contains the following substances.

For Examiner's Use

- phosphorus, P<sub>4</sub>
- potassium chlorate(V), KClO<sub>3</sub>
- sulphur, S
- a hydrocarbon wax

(a)	the Pho	friction between the match head and the striking surface generates enough heat for phosphorus to burn. sphorus burns to form phosphorus(V) oxide. This oxide is covalently bonded with a ecular structure.
	(i)	What is the molecular formula of phosphorus(V) oxide?
		[1]
	(ii)	Suggest <b>one</b> physical and <b>one</b> chemical property of phosphorus(V) oxide.
		physical property
		chemical property
		[2]
(b)	dec	heat from the combustion of phosphorus provides enough energy for the omposition of potassium chlorate( $V$ ) to oxygen and potassium chloride. estruct the equation for the decomposition of potassium chlorate( $V$ ).
		[2]
(c)		sulphur on the match head ignites. e an equation to show the combustion of sulphur.
		[1]
(d)	One	ally the wax on the match head begins to combust. compound in the wax has the formula $C_{18}H_{38}$ . which class of hydrocarbons does this compound belong? Explain your answer.

[Total: 7]

<b>A6</b>	Sulphur dioxide,	SO <sub>2</sub> , and	nitrogen	dioxide,	NO <sub>2</sub> ,	are	both	atmospheric	pollutants	formed
	during the combi	ustion of co	oal at a po	ower stat	ion.					

(a) (i) State another source of sulphur dioxide as an atmospheric pollutant.

\_\_\_\_\_[1]

(ii) State another source of nitrogen dioxide as an atmospheric pollutant.

\_\_\_\_\_[1]

**(b)** Nitrogen dioxide and sulphur dioxide both cause acid rain. They are removed from the flue gases released from the power station by reaction with moist calcium carbonate in a process called flue gas desulphurisation.

Calcium carbonate reacts with sulphur dioxide to make a solid called calcium sulphite and a gas.

(i) What is the name of this gas?

.....[1]

(ii) Nitrogen dioxide reacts with calcium carbonate to make a solid. Suggest the name of this solid.

[1]

(iii) Describe one environmental effect of acid rain.

.....[1]

**(c)** Sulphur dioxide and nitrogen dioxide react together as shown in the equation.

$$SO_2 + NO_2 \rightarrow SO_3 + NO$$
  $\Delta H = +43 \text{ kJ/mol}$ 

Draw an energy profile diagram for this reaction.

Indicate both the enthalpy change and the activation energy on your diagram.



[Total: 8]

[3]

## **Section B**

Answer **three** questions from this section.

For Examiner's Use

The total mark for this section is 30.

- **B7** This question is about the chemistry of chlorine and some of its compounds.
  - (a) Describe, with the aid of an ionic equation, the reaction of chlorine with aqueous potassium bromide. Explain why this reaction involves the reduction of chlorine. [3]
  - (b) Magnesium reacts with chlorine to form magnesium chloride.
     Draw diagrams to show the electronic structures and charges of both ions present in magnesium chloride.
  - (c) Silver chloride is an insoluble salt.

    Outline the preparation of pure, dry silver chloride, starting from solid silver nitrate. [4]
  - (d) State **one** environmental problem associated with the molecule  $C_2F_3Cl_3$ . [1]

[Total: 10]

**B8** Crude oil is a raw material which is processed in an oil refinery. Two of the processes used are fractional distillation and cracking.

For Examiner's Use

The table shows the percentage by mass of some different fractions in crude oil. The table also shows the demand for each fraction expressed as a percentage.

fraction	number of carbon atoms per molecule	percentage in crude oil	percentage needed by the oil refinery to supply demand		
petroleum gases	1 - 4	4%	11%		
gasoline	5 - 9	11%	22%		
kerosene	10 - 14	12%	20%		
gas oil	14 - 20	18%	15%		
waxes and bitumen	over 20	23%	4%		

- (a) The variation in which physical property is used to separate crude oil by fractional distillation?
- (b) (i) Define the term cracking.

[2]

- (ii) Use information from the table to explain how cracking helps an oil refinery match the supply of gasoline with the demand for gasoline. [2]
- (c) The hydrocarbon  $C_{15}H_{32}$  can be cracked to make propene and one other hydrocarbon.
  - (i) Draw the structure of propene.

[1]

(ii) Write an equation for this reaction.

[1]

[1]

- **(d)** Propene is used to make alcohols and poly(propene).
  - (i) Describe how propene can be converted into an alcohol and draw the structure of this alcohol. [2]
  - (ii) Draw the structure of poly(propene) showing at least two repeat units.

[Total: 10]

© UCLES 2008 5070/02/M/J/08

**B9** Dilute ethanoic acid and dilute hydrochloric acid both react with magnesium ribbon to form hydrogen.

For Examiner's Use

(a) Give the formula of one ion found in both of these dilute acids.

- [1]
- (b) Magnesium ribbon reacts with hydrochloric acid as shown in the equation.

$$Mg + 2HCl \rightarrow MgCl_2 + H_2$$

A 0.24g sample of magnesium ribbon is added to 5.0 cm<sup>3</sup> of 2.0 mol/dm<sup>3</sup> hydrochloric acid.

- (i) Which reactant, magnesium or hydrochloric acid, is in excess? Use calculations to explain your answer. [2]
- (ii) Calculate the maximum mass of magnesium chloride that can be formed in this reaction. [2]
- (iii) A 0.24g sample of magnesium ribbon is added to 5.0 cm³ of 2.0 mol/dm³ ethanoic acid.
   Explain why this reaction forms the same volume of hydrogen but takes place much more slowly than the reaction of the same mass of magnesium with 5.0 cm³ of 2.0 mol/dm³ hydrochloric acid.
- (c) (i) Write an equation for the reaction between dilute ethanoic acid and sodium carbonate. [1]
  - (ii) What observations would be made during this reaction? [1]

[Total: 10]

B10 Brass is an alloy of zinc and copper.

For Examiner's Use

- (a) Describe, with the aid of a labelled diagram, the structure of a metal such as copper. [2]
- (b) Explain, in terms of their structures, why both zinc and copper are good conductors of electricity. [1]
- (c) A 1.2 g sample of powdered brass was analysed by reaction with excess dilute sulphuric acid.

The zinc reacts as shown in the equation to form 0.072 dm³ of hydrogen measured at room temperature and pressure.

$$Zn + 2H^+ \rightarrow Zn^{2+} + H_2$$

- (i) Suggest why brass was used in a powdered rather than lump form. [1]
- (ii) Calculate the mass of zinc in the sample of brass. [2]
- (iii) Calculate the percentage of zinc in the sample of brass. [1]
- (d) Describe how aqueous ammonia can be used to show that only the zinc in the sample reacted with the acid. [3]

[Total: 10]

e clearly shown.	

L

For Examiner
Use
1

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

DATA SHEET

		0	4	Helium C	7		Neon 10	40	Ā	Argon 18	8	궃	Krypton 36	131	Xe	Xenon 54	222	Ru	Radon 86				175	Γn	Lutetium
		NII VII			ç	₽ <b>L</b>	Fluorine 9	35.5	73	Chlorine 17	80	ā	Bromine 35	127	П	lodine 53	210	Ą	Astatine 85				173	Υp	Ytterbium
		IN			4	₽ <b>O</b>	Oxygen 8	32	တ		62	Se	Selenium 34	128	<u>e</u>	Tellurium 52	209	Po	Polonium 84				169	Ę	Thulium
		^			7	<b>₹ Z</b>	Nitrogen 7	31	凸	Phosphorus 15	75	As		122	Sb	Antimony 51	209	Ξ	Bismuth 83				167	й	Erbium
		$\geq$			ć	2 <b>U</b>	Carbon 6	28	Si	Silicon 14	73	Ge	Germanium 32	119	Sn	Tin 50	207	Pb	Lead 82				165	운	Holmium
					7	= <b>m</b>	Boron 5	27	ΡΙ	Aluminium 13	02	Ga	Gallium 31	115	In	Indium 49	204	11	Thallium 81				162	۵	Dysprosium
ts												Zu	Zinc 30	112	ဦ	Cadmium 48	201	Нg	Mercury 80				159	욘	Terbium
The Periodic Table of the Elements											49	Cn	Copper 29	108	Ag	Silver 47	197	Au	Gold 79				157	gq	Gadolinium
e of the	Group										29	Z	Nickel 28	106	Pd	Palladium 46	195	Ŧ	Platinum 78				152	En	Europium
dic Tabl	Gre										29	ပိ	Cobalt 27	103	Rh	Rhodium 45	192	ľ	Iridium 77				150	Sm	Samarium
he Perio			- 3	Î	-						26	Ьe	Iron 26	101	Ru	Ruthenium 44	190	Os	Osmium 76				147	Pm	Promethium
T											55	M	Manganese 25		ည	Technetium 43	186	Re	Rhenium 75				144	N	Neodymium
											52	ပ်	Chromium 24	96	Mo	Molybdenum 42	184	>	Tungsten 74				141	ፈ	Praseodymium
											51	>	Vanadium 23	63	qN	Niobium 41	181	<u>n</u>	Tantalum 73				140	ဝီ	Cerium
											48	F	Titanium 22	91	Zr	Zirconium 40	178	Ξ	Hafnium 72				7		
											45	လွ	Scandium 21	68	>	Yttrium 39	139	Гa	Lanthanum 57 *	227	Ac	Actinium 89 †	opripo	2000	SGIRO
		=			d	» e	Beryllium 4	24	Mg	Magnesium 12	40	Sa	Calcium 20	88	Š	Strontium 38	137	Ва	Barium 56	226	Ra	Radium 88	* 58_71 Lanthanoid series	+ OO +OO Actinoid corion	O ACIII IOIO
		_			1	\ <u> </u>	Lithium 3	23	Na	Sodium 11	39	¥	Potassium 19	85	Rb	Rubidium 37	133	Cs	Caesium 55	223	ŗ	Francium 87	* 58_71	, ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ; ;	30 I OS I
CLES	S 200	18										507	0/02	/M/.	I/N8	2									

	4	7	144	4 47	740	450	457	024	460		467	760	473	175
anoid series	140	141	144	14/	OGI.	761	/61	601	791		/91	691	5/1	6/1
oid sprips	ပီ	Ą	Š	Pm	Sm	Ш	gg	Q L	ک	웃	ш	E	Υp	3
	Cerium 58	Praseodymium 59	Neodymium 60	Promethium 61	Samarium 62	Europium 63	Gadolinium 64	Terbium 65	Dysprosium 66	67	Erbium 68	Thulium 69	Ytterbium 70	Lutetium 71
a = relative atomic mass	232	231	238	237	244	243	247	247	251	252	257	258	259	260
X = atomic symbol	۲	Ра	<b>-</b>	d N	Pu	Am	Cu	æ	చ	Es	Fm	Md	8	בֿ
b = atomic (proton) number	Thorium 90	Protactinium 91	Uranium 92	Neptunium 93	Plutonium 94	Americium 95	Curium 96	Berkelium 97	Californium 98	Einsteinium 99	Fermium 100	Mendelevium 101	Nobelium 102	Lawrencium 103

Key

The volume of one mole of any gas is 24dm3 at room temperature and pressure (r.t.p.).