CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Ordinary Level

MARK SCHEME for the October/November 2012 series

5070 CHEMISTRY

5070/22 Paper 2 (Theory), maximum raw mark 75

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2012 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



	Pa	ge 2		Mark Scheme	Syllabus	Paper
				GCE O LEVEL – October/November 2012	5070	22
A 1	(a)	(sul	ostan	ce containing) two or more elements bonded / joine	ed	[1]
	(b)	(i)	carb	on dioxide / CO ₂		[1]
		(ii)	zinc	oxide / ZnO		[1]
		(iii)	calci	ium carbonate / CaCO₃		[1]
		(iv)	carb	on dioxide / CO ₂		[1]
		(v)	meth	nane / CH ₄		[1]
		(vi)	carb	on monoxide / CO		[1]
	(c)	one	pair	of electrons between each H and O; (1)		
		rest	of st	ructure is correct; (1)		[2]
						[Total: 9]
A2	(a)	(i)	lead	< iron < zinc < magnesium		[1]
		(ii)	Fe ₂ C	D_3 + 3Zn \rightarrow 3ZnO + 2Fe		[1]
	(b)	(i)	(forn	ns an) oxide layer / has a coat of oxide; (1)		
				ch is strongly fixed to the surface / which is not eacactive; (1)	sily removed / wh	nich is [2]
		(ii)	low	density		[1]
		(iii)	proto	ons = 13 and neutrons = 14		[1]
						[Total: 6]

Page 3	Mark Scheme	Syllabus	Paper
	GCE O LEVEL – October/November 2012	5070	22

A3 NOTE: for parts A3a(i) and A3a(ii) answers must be comparative

(a) (i) speed increases with increase in bromine concentration (no mark alone) because

(bromine) molecules closer together / more (bromine) molecules (in a given volume) / more (bromine) particles (in a given volume) / more crowded molecules; (1)

therefore frequency of collisions greater /more particles collide per second / greater chance of collisions / collide more often; (1)

[2]

(ii) increasing temperature increases rate (no mark alone) because

particles move more rapidly / particles have more energy; (1)

therefore more energetic collisions / more effective collisions / more successful collisions / more vigorous collisions; (1)

NOTE: more particles have energy greater than activation energy = 2 marks

[2]

(iii) measure colour of the solution / bromine (over time) / use a colorimeter / measure absorbance / measure how much light goes through the solution / measure (electrical) conductivity

[1]

(b) (i) Fe
$$\to$$
 Fe²⁺ + 2e⁻ (1)

$$Br_2 + 2e^- \rightarrow 2Br^- (1)$$
 [2]

(ii) reactants on the left and products on the right and reactant level above product level; (1)

 ΔH correctly labelled with arrow going downwards; (1)

activation energy correctly labelled with arrow / line going upwards or double-headed arrow; (1)

[Total: 10]

[3]

Page 4	Mark Scheme	Syllabus	Paper
	GCE O LEVEL – October/November 2012	5070	22

A4 (a) carbon dioxide and water (required); (1)

(in presence of) sunlight / chlorophyll; (1)

to form glucose / $C_6H_{12}O_6$ / sugars / carbohydrate; (1) [3]

(b) (i) calcium ethanoate

(ii) boiling point [1]

(iii) C = 54.5/12 H = 9.1/1 O = 36.4/16 or

4.54 9.1 2.275 / 2.28 (1)

ratio = 2 4 1 (1) [2]

(c) (i) formula completed correctly e.g. $- OCH_2CH_3$ [1]

(ii) solvent / flavouring / perfumes / making polyesters / making terylene / plasticisers / making fuels (transesterification) / nail varnish remover [1]

[Total: 9]

[1]

				GCE O LEVEL – October/November 2012	5070	22
A 5	(a)	(i)	evap	porates easily / easily form a gas		[1
		(ii)	by h	eating / high temperature		[1
		(iii)		urities remain as solids / impurities do not evonyl evaporates / nickel reacts and leaves impurit		e nickel [1
	(b)	4				[1
	(c)	two	elect	trodes dipping into liquid and power pack or batter	y; (1)	
		(pu	re) ni	ckel and impure nickel electrodes labelled; (1)		
		imp	oure n	ickel is the anode / + electrode and pure nickel is	the cathode / - ele	ctrode; (1)
		ele	ctrolyt	te labelled as nickel salt / named nickel salt / aque	ous nickel compou	und; (1) [4
	(d)	•		e from: s heat / conducts electricity (1)		
		ma	lleable	e / can be hammered into shape / can be bent into	shapes (1)	
		duc	ctile / d	can be stretched (1)		
		shii	ny / lu	strous (1) IGNORE: silvery		[3

Mark Scheme

Syllabus

Paper

[Total: 11]

Page 5

Page 6	Mark Scheme	Syllabus	Paper
	GCE O LEVEL – October/November 2012	5070	22

B6 (a) (i) chlorine gains electrons, so is reduction; (1)

- (iii) iodine is less reactive (than bromine) ORA iodine is lower in the reactivity series (than bromine) [1]
- (b) C and D because they have low boiling points/C and D because they do not conduct (when molten)[1]

(c)
$$Cl_2 + 2NaOH \rightarrow NaClO + NaCl + H_2O$$
 [1]

(d) (i)
$$0.05 \text{ (mol dm}^{-3}\text{)}$$

(ii) mol thiosulfate =
$$0.05 \times 23.6/1000 / 1.18 \times 10^{-3}$$
 (mol); (1)

mol iodine =
$$5.9 \times 10^{-4}$$
 (mol); (1)

concentration of iodine =
$$(5.9 \times 10^{-4} \times 1000 / 12.5) = 0.0472 \text{ (mol dm}^{-3}) \text{ (1)}$$
 (mark is for correct answer) [3]

[Total: 10]

Page 7	Mark Scheme	Syllabus	Paper
	GCE O LEVEL – October/November 2012	5070	22

B7 (a) (i) (both have) tetrahedral arrangement of atoms / (both have) hexagonal arrangement of atoms; (1) (both are) giant structures / giant molecular (structures) / macromolecules / [2] covalent lattices; (1) (ii) many (covalent) bonds / giant structure / macromolecule / all atoms joined together / network of bonds / lattice; takes a lot of energy to break bonds / hard to break bonds / high temperature needed to break bonds / bonds are strong; [2] (iii) no free electrons / no delocalised electrons / no sea of electrons / all electrons in covalent bonds / electrons can't move / electrons in fixed positions; [1] (b) (i) idea of random movement of molecules or particles / movement of molecules or particles in any direction; NOTE: answer must refer to particles, of any kind [1] (ii) they have different masses / they have different sizes / hydrogen (ion) is lighter / hydrogen (ion) is smaller [1]

charges correct Na⁺ and O²⁻; (1)

2 sodium ions and 1 oxide ion / Na₂O / ratio of 2 Na to 1 O from diagram of covalent structure; (1)

[3]

Page 8	Mark Scheme	Syllabus	Paper
	GCE O LEVEL – October/November 2012	5070	22

B8 (a) improve (crop) growth / improve (crop) yield / increase crop (growth) / increase crop (yield) / bigger crop (growth) / better crop (yield)

[1]

(b) so that the roots can absorb them / so the plant can absorb them

[1]

(c) (i) 2N = 28; (1)

RFM of ammonium nitrate = 80; (1)

% by mass = 35%; (1)

[3]

(c) (ii) add sulfuric acid (to aqueous ammonia); (1)

titration; (1)

use of indicator then repeat without indicator; (1)

heat (solution obtained) to crystallisation point / evaporate some of the water (from the solution) / heat (solution) then leave (solution) to cool / leave (solution) to crystallise / solution concentrated by heating (1)

[4]

[1]

(d) 3-

[Total: 10]

Page 9	Mark Scheme	Syllabus	Paper
	GCE O LEVEL – October/November 2012	5070	22

B9 (a) (i) sulfur dioxide / hydrogen peroxide

[1]

(ii) kills bacteria

[1]

(b) (i) correct structure with

two or more units and single bonds between carbon atoms; (1)

continuation bonds present; (1)

[2]

(ii) bromine water / (aqueous) bromine / bromine; (1)

turns colourless / decolourised; (1)

[2]

(c) (i) correct formula for ethanoate ion showing all atoms and bonds including negative charge on the single bonded oxygen

[1]

(ii) $CH_3COOH + OH^- \rightarrow CH_3COO^- + H_2O$

[1]

(d) (i) (hydroxide reacts with ammonium salts) to form ammonia

[1]

(ii) $OH^- + NH_4^+ \rightarrow NH_3 + H_2O$

[1]

[Total: 10]