


# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Ordinary Level

| CANDIDATE<br>NAME |  |  |                     |  |  |
|-------------------|--|--|---------------------|--|--|
| CENTRE<br>NUMBER  |  |  | CANDIDATE<br>NUMBER |  |  |



CHEMISTRY 5070/21

Paper 2 Theory May/June 2013

Candidates answer on the Question Paper.

No Additional Materials are required.

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

#### **Section A**

Answer all questions.

Write your answers in the spaces provided in the Question Paper.

#### **Section B**

Answer any three questions.

Write your answers in the spaces provided in the Question Paper.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 20.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

This document consists of 17 printed pages and 3 blank pages.



1 hour 30 minutes

### **Section A**

For Examiner's Use

[Total: 6]

Answer all the questions in this section in the spaces provided.

The total mark for this section is 45.

A1 Choose from the following compounds to answer the questions below.

#### butane

calcium carbonate
carbon dioxide
copper(II) nitrate
iron(II) hydroxide
iron(III) hydroxide
propene
sodium chloride
sulfur dioxide
sulfuric acid

Each compound can be used once, more than once or not at all.

Name a compound which

| (a) | is a green solid,  |
|-----|--|
|     | [1]  |
| (b) | is a saturated hydrocarbon,  |
|     | [1]  |
| (c) | has a molecule with only 9 atoms,  |
|     | [1]  |
| (d) | can be used to reduce the acidity in lakes,                                |
|     | [1]  |
| (e) | will turn aqueous acidified potassium dichromate(VI) from orange to green, |
|     | [1]  |
| (f) | can be electrolysed in aqueous solution to form two gases.                 |
|     | [1]  |
|     |  |

For Examiner's Use

[Total: 9]


| <b>A2</b> | Pho | otosynthesis helps to maintain the percentage of oxygen in air. |   |  |  |  |  |  |  |
|-----------|-----|---|---|--|--|--|--|--|--|
|           | (a) |   | at is the percentage, by volume, of oxygen in dry air?  |  |  |  |  |  |  |
|           | (b) | In a  | ddition to releasing oxygen, photosynthesis produces glucose, $C_6H_{12}O_6$ .  e the overall equation that represents photosynthesis.  |  |  |  |  |  |  |
|           | (c) | Des   | cribe the essential conditions needed for photosynthesis.   |  |  |  |  |  |  |
|           |     |   | [2]   |  |  |  |  |  |  |
|           | (d) | Pho   | tosynthesis is an endothermic reaction.   |  |  |  |  |  |  |
|           | (4) | (i)   | Explain, in terms of the energy changes that occur during bond breaking and bond making, why photosynthesis is an endothermic reaction. |  |  |  |  |  |  |
|           |     |   |   |  |  |  |  |  |  |
|           |     |   | [2]   |  |  |  |  |  |  |
|           |     | (ii)  | Complete the energy profile diagram for photosynthesis.   |  |  |  |  |  |  |
|           |     |   | On your diagram label the<br>• products,<br>• enthalpy change for the reaction, $\Delta H$ ,<br>• activation energy, $E_{\rm a}$ .      |  |  |  |  |  |  |
|           |     |   |   |  |  |  |  |  |  |
|           |     |   | energy reactants  |  |  |  |  |  |  |
|           |     |   | progress of reaction  |  |  |  |  |  |  |
|           |     |   | [3]   |  |  |  |  |  |  |

A3 Salts are often made by the neutralisation of bases.

For Examiner's Use

(a) Aqueous potassium hydroxide, of concentration 0.150 mol/dm³, is added to 25.0 cm³ of sulfuric acid in a flask.

The graph shows how the pH of the liquid in the flask changes as aqueous potassium hydroxide is added to it.



(i) Construct the equation for the complete neutralisation of sulfuric acid by potassium hydroxide.

.....[1]

(ii) Use the graph to deduce the volume of aqueous potassium hydroxide required to neutralise 25.0 cm<sup>3</sup> of sulfuric acid.

.....[1]

|     | (iii) | Use your answers to (i) and (ii) to calculate the concentration of sulfuric acid.   | For<br>Examiner's<br>Use |
|-----|-------|---|--------------------------|
| (b) |       | concentration of sulfuric acid = mol/dm <sup>3</sup> [3] scribe the essential experimental details for preparing a pure sample of zinc nitrate stals from zinc oxide. |                          |
|     |       | stats from Zine oxide.  |                          |
|     |       |   |                          |
|     |       |   |                          |
|     |       | [4]   |                          |
|     |       | [Total: 9]  |                          |

# **BLANK PAGE**

A4 The table shows the number of electrons, neutrons and protons in seven different particles.

For Examiner's Use

| particle | number of |          |         |  |  |
|----------|-----------|----------|---------|--|--|
| particle | electrons | neutrons | protons |  |  |
| Α        | 12        | 12       | 12      |  |  |
| В        | 15        | 16       | 15      |  |  |
| С        | 17        | 18       | 17      |  |  |
| D        | 17        | 20       | 17      |  |  |
| E        | 18        | 16       | 16      |  |  |
| F        | 18        | 22       | 18      |  |  |
| G        | 18        | 20       | 20      |  |  |

| (a) | What is the nucleon number for <b>F</b> ?         |            |
|-----|---|------------|
|     |   | [1]        |
| (b) | Explain why A is a neutral particle.              |            |
|     |   |            |
|     |   |            |
|     |   | [2]        |
| (c) | Which particles are isotopes of the same element? |            |
|     |   | [1]        |
| (d) | What is the charge on <b>E</b> ?                  |            |
|     |   | [1]        |
| (e) | Which particles have the same relative mass?      |            |
|     |   | [1]        |
|     |   | [Total: 6] |

**A5** Analysis of compound **X** shows it has the following composition.

| For        |
|------------|
| Examiner's |
| Use        |

[3]

| element  | percentage by mass |
|----------|--------------------|
| nitrogen | 11.1               |
| hydrogen | 3.20               |
| chromium | 41.3               |
| oxygen   | 44.4               |

(a) Show that  ${\bf X}$  has the formula  ${\bf N_2H_8Cr_2O_7}.$ 

| (b) | An a | aqueous solution of <b>X</b> is orange.   |    |
|-----|------|---|----|
|     | Sug  | gest which element in <b>X</b> is responsible for the orange colour.  |    |
|     |      | [   | 1] |
| (c) | An a | acidified aqueous solution of ${f X}$ reacts with aqueous potassium iodide to form iodine                                   | е. |
|     | Stat | te and explain what you can conclude about the chemical nature of <b>X</b> .  |    |
|     |      |   |    |
|     |      |   |    |
|     |      | [2  | 2] |
| (d) |      | leous sodium hydroxide is added to solid ${\bf X}$ and the mixture is warmed. A gas the s moist red litmus blue is evolved. | at |
|     | (i)  | Give the formula of the positive ion present in <b>X</b> .  |    |
|     |      | [   | 1] |
|     | (ii) | Suggest the formula of the other ion present in X.  |    |
|     |      | ſ·  | 11 |

| (e) | When solid <b>X</b> is heated only $Cr_2O_3$ , water and gas <b>Z</b> are formed. | For               |
|-----|---|-------------------|
|     | Name gas <b>Z</b> .   | Examiner's<br>Use |
|     | [1]   |                   |
|     | [Total: 9]  |                   |

Question A6 starts on page 10.

For Examiner's Use

| <b>A6</b> | Potassium is in Group I and chlorine is in Group VII of the Periodic Table. |              |   |  |
|-----------|---|--------------|---|--|
|           | Pota<br>Cl <sub>2</sub> (   |              | m forms an oxide with the formula ${\rm K_2O}$ and chlorine forms an oxide with the formula   |  |
|           | (a)   | (i)          | Draw a 'dot-and-cross' diagram for ${\rm C}\it{l}_{\rm 2}{\rm O}.$  |  |
|           |   |              | You only need to draw the outer shell electrons.  |  |
|           |   |              |   |  |
|           |   |              |   |  |
|           |   |              | [1]   |  |
|           |   | (ii)         | Explain, using ideas about structure and bonding, why $\mathrm{C}\mathit{l}_2\mathrm{O}$ has a low melting point.   |  |
|           |   |              |   |  |
|           |   |              |   |  |
|           |   |              | [2]   |  |
|           | (b)   |              | w diagrams to show the electronic structures and charges of both ions present in assium oxide.  |  |
|           |   |              |   |  |
|           |   |              |   |  |
|           |   |              |   |  |
|           |   |              | [2]   |  |
|           | (c)   | Chlo<br>wate | prine forms another oxide $\mathrm{C}l_2\mathrm{O}_7$ . One mole of this oxide reacts with one mole of er to make two moles of an acid and no other products. |  |
|           |   | Con          | struct the equation for this reaction.  |  |
|           |   |              | [1]   |  |
|           |   |              | [Total: 6]  |  |
|           |   |              |   |  |

## **Section B**

For Examiner's Use

Answer three questions from this section in the spaces provided.

The total mark for this section is 30.

| 37 | Mal | achit | e is an ore of copper. The formula of malachite is ${\rm CuCO_3.Cu(OH)_2.}$  |
|----|-----|-------|--|
|    | Mal | achit | e reacts as though it is a mixture of copper(II) carbonate and copper(II) hydroxide.   |
|    |     |       | sample of malachite is added to excess dilute hydrochloric acid, $HCl(aq)$ . The carbon ormed is collected and has a volume of $96\mathrm{cm}^3$ at room temperature and pressure. |
|    | (a) | Wha   | at would you observe when malachite reacts with HCl(aq)?   |
|    |     |       |  |
|    |     |       | [2]  |
|    | (b) | Con   | struct the equation for the reaction between malachite and $HCl(aq)$ .   |
|    |     |       | [2]  |
|    | (c) | Cald  | culate the mass of carbonate ion, ${\rm CO_3}^{2-}$ , in the sample of malachite.  |
|    |     |       |  |
|    |     |       |  |
|    |     |       |  |
|    |     |       |  |
|    |     |       |  |
|    |     |       | mass of $CO_3^{2-} = \dots g [3]$  |
|    | (d) | Cop   | per is extracted from malachite by heating with carbon.  |
|    |     | (i)   | Construct an equation for the reduction of malachite by carbon.  |
|    |     |       | [2]  |
|    |     | (ii)  | Malachite is a finite resource. Give one <b>other</b> reason why copper should be recycled.  |
|    |     |       |  |

[Total: 10]

......[1]

**B8** Carboxylic acids are a homologous series of organic compounds.

For Examiner's Use

The table shows information about some carboxylic acids.

| carboxylic acid   | formula   | melting point/°C | boiling point/°C |
|-------------------|---|------------------|------------------|
| methanoic acid    | HCO <sub>2</sub> H                                | 8                | 100              |
| ethanoic acid     | CH <sub>3</sub> CO <sub>2</sub> H                 | 17               | 118              |
|                   | C <sub>2</sub> H <sub>5</sub> CO <sub>2</sub> H   | -22              | 141              |
| butanoic acid     | C <sub>3</sub> H <sub>7</sub> CO <sub>2</sub> H   |                  |                  |
| hexadecanoic acid | C <sub>15</sub> H <sub>31</sub> CO <sub>2</sub> H | 63               | 269              |

| (a) | What is meant by the term <i>homologous series</i> ?  |
|-----|---|
|     |   |
|     |   |
|     | [2]   |
| (b) | Name the carboxylic acid with the formula $\mathrm{C_2H_5CO_2H}$ .  |
|     | [1]   |
| (c) | Deduce the general formula for a carboxylic acid.   |
|     | [1]   |
| (d) | It is more difficult to estimate the melting point of butanoic acid than its boiling point. Use the data in the table to explain why. |
|     |   |
|     | [1]   |
| (e) | When warmed in the presence of concentrated sulfuric acid, butanoic acid reacts with ethanol to make an ester.                        |
|     | Name and draw the structure, showing all the atoms and all the bonds, of this ester.  |
|     | name  |
|     | structure   |

| (f) | Hex  | adecanoic acid, C <sub>15</sub> H <sub>31</sub> CO <sub>2</sub> H, is a weak acid.  | For     |
|-----|------|---|---------|
|     | (i)  | Write an equation to show the dissociation of hexadecanoic acid when dissolved in water. Use the equation to explain the meaning of the term weak acid. | Use Use |
|     |      |   |         |
|     |      | [2]   |         |
|     | (ii) | What is the formula of the salt formed when hexadecanoic acid reacts with aqueous sodium hydroxide?   |         |
|     |      | [1]   |         |
|     |      | [Total: 10]   |         |

| B9 | Ethanol is | manufactured b    | v the h   | vdration o  | f ethene |
|----|------------|-------------------|-----------|-------------|----------|
|    |            | THAILAILAOLAICA D | y 1110 11 | y aration o |          |

| For        |
|------------|
| Examiner's |
| llse       |

$$C_2H_4(g) + H_2O(g) \rightleftharpoons C_2H_5OH(g)$$
  $\Delta H = -45 \text{ kJ/mol}$ 

This reaction is exothermic.

The reaction is normally carried out at a pressure of 70 atmospheres and a temperature of  $300\,^{\circ}\text{C}$ .

| 000 | ٥.      |   |
|-----|---------|---|
| (a) | The     | reaction is carried out at 70 atmospheres pressure and at <b>600 °C</b> rather than 300 °C. |
|     | Pred    | dict and explain the effect of raising the temperature on                                   |
|     | (i)     | the rate of reaction,   |
|     |         |   |
|     |         |   |
|     |         | [2]   |
|     | (ii)    | the position of equilibrium.  |
|     |         |   |
|     |         |   |
|     |         | [2]   |
| (b) | The 300 | reaction is carried out at <b>20 atmospheres</b> rather than 70 atmospheres, and at °C.     |
|     | Pred    | dict and explain the effect of decreasing the pressure on                                   |
|     | (i)     | the rate of reaction,   |
|     |         |   |
|     |         |   |
|     |         | [2]   |
|     | (ii)    | the position of equilibrium.  |
|     |         |   |
|     |         |   |
|     |         | [2]   |

| (C) | Calculate the energy released when 10 moles of ethanol are formed. | For<br>Examiner's<br>Use |
|-----|--|--------------------------|
|     | energy released =kJ [1]  |                          |
| (d) | The hydration of ethene uses an acid catalyst.                     |                          |
|     | Explain how a catalyst can increase the rate of reaction.          |                          |
|     |  |                          |
|     | [1]  |                          |
|     | [Total: 10]  |                          |

**B10** Aqueous silver nitrate can be electrolysed using inert electrodes. Solid silver is formed on the cathode (negative electrode).

For Examiner's Use

The table shows how the mass of silver formed is affected by four factors.

| temperature<br>of solution<br>/°C | duration of<br>electrolysis<br>/seconds | current passed<br>through<br>solution/amps | concentration of<br>solution<br>/mol/dm <sup>3</sup> | mass of silver<br>formed<br>/g |
|-----------------------------------|---|--|--|--------------------------------|
| 25                                | 100                                     | 9.65                                       | 1.0  | 0.108                          |
| 30                                | 100                                     | 9.65                                       | 1.0  | 0.108                          |
| 25                                | 100                                     | 9.65                                       | 0.5  | 0.108                          |
| 25                                | 200                                     | 9.65                                       | 0.5  | 0.216                          |
| 25                                | 100                                     | 19.3                                       | 1.0  | 0.216                          |

| (a) | The  | electrode reaction at the cathode is reduction.  |
|-----|------|--|
|     | (i)  | Construct the equation for the reaction which occurs at the cathode.                   |
|     |      | [1]  |
|     | (ii) | Explain why this reaction is reduction.  |
|     |      |  |
|     |      | [1]  |
| (b) | Stat | te how each of the following factors affects the mass of silver formed at the cathode. |
|     | tem  | perature of solution   |
|     |      |  |
|     |      |  |
|     | dura | ation of electrolysis  |
|     |      |  |
|     |      |  |
|     | curr | rent used  |
|     |      |  |
|     |      |  |
|     | con  | centration of solution   |
|     |      |  |
|     |      |  |

[4]

| (C) | Explain why aqueous sliver nitrate can be electrolysed but solid sliver nitrate cannot.             | For<br>Examiner's<br>Use |
|-----|---|--------------------------|
|     |   |                          |
|     | [2]   |                          |
| (d) | Aqueous silver nitrate reacts with dilute hydrochloric acid to form a white precipitate.            |                          |
|     | Construct the ionic equation, including state symbols, for the formation of this white precipitate. |                          |
|     | [2]   |                          |
|     | [Total: 10]   |                          |

# **BLANK PAGE**

## **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

DATA SHEET
The Periodic Table of the Elements

|                                   |       |     |   |          |   |    |    |                |      |    |                  |    |        |                 |      |          |                  |     |          |                   | 1   |    |                |                           |               |                          |
|-----------------------------------|-------|-----|---|----------|---|----|----|----------------|------|----|------------------|----|--------|-----------------|------|----------|------------------|-----|----------|-------------------|-----|----|----------------|---------------------------|---------------|--------------------------|
|                                   |       | 0   | 4 | ¥ ‡      | 2 | 20 | Ne | Neon<br>10     | 40   | Ar | Argon<br>18      | 84 | ¥      | Krypton<br>36   | 131  | Xe       | Xenon<br>54      | 222 | R        | Radon<br>86       |     |    |                | 175                       | Ľ             | Lutetium                 |
|                                   |       | IIΛ |   |          |   | 19 | ш  | Fluorine<br>9  | 35.5 | CI | Chlorine<br>17   | 80 | ģ      | Bromine<br>35   | 127  | Ι        | lodine<br>53     | 210 | Αt       | Astatine<br>85    |     |    |                | 173                       | Υb            | Ytterbium                |
|                                   |       | ΙΛ  |   |          |   | 16 | 0  | Oxygen<br>8    | 32   | S  |                  | 79 | Se     | Selenium<br>34  | 128  | <u>a</u> | Tellurium<br>52  | 209 | Ъ        | Polonium<br>84    |     |    |                | 169                       | E             | Thulium                  |
|                                   |       | >   |   |          |   | 14 | z  | Nitrogen<br>7  | 31   | ۵  | Phosphorus<br>15 | 75 | As     | Arsenic<br>33   | 122  | Sb       | Antimony<br>51   | 209 | <u>.</u> | Bismuth<br>83     |     |    |                | 167                       | ш             | Erbium                   |
|                                   |       | 2   |   |          |   | 12 | ပ  | Carbon<br>6    | 28   | Si | _                | 73 | Ge     | Germanium<br>32 | 119  | Sn       | Tin<br>50        | 207 | Pb       | Lead<br>82        |     |    |                | 165                       | 운             | Holmium                  |
|                                   |       | ≡   |   |          |   | Ξ  | Δ  | Boron<br>5     | 27   | ΝI | Aluminium<br>13  | 70 | Са     | Gallium<br>31   | 115  | In       | Indium<br>49     | 204 | 11       | Thallium<br>81    |     |    |                | 162                       | ۵             | Dysprosium               |
| S                                 |       |     |   |          | • |    |    |                |      |    |                  | 65 | Zu     | Zinc<br>30      | 112  | ပ္ပ      | Cadmium<br>48    | 201 | Hg       | Mercury<br>80     |     |    |                | 159                       | Д             | Terbium                  |
| Element                           |       |     |   |          |   |    |    |                |      |    |                  | 64 | D<br>C | Copper<br>29    | 108  | Ag       | Silver<br>47     | 197 | Au       | Gold<br>79        |     |    |                | 157                       | gg            | Gadolinium               |
| he Periodic Table of the Elements | Group |     |   |          |   |    |    |                |      |    |                  | 59 | Z      | Nickel<br>28    | 106  | Pd       | Palladium<br>46  | 195 | Ŧ        | Platinum<br>78    |     |    |                | 152                       | Ш             | Europium                 |
| dic Table                         | Gro   |     |   |          |   |    |    |                |      |    |                  | 59 | ပိ     | Cobalt<br>27    | 103  | R        | Rhodium<br>45    | 192 | <u> </u> | Iridium<br>77     |     |    |                | 150                       | Sm            | Samarium                 |
| ne Perio                          |       |     | - | <b>T</b> | 1 |    |    |                |      |    |                  | 56 | Ъ      | Iron<br>26      | 101  | Bu       | Ruthenium<br>44  | 190 | SO.      | Osmium<br>76      |     |    |                | 147                       | Pm            | Promethium               |
| F                                 |       |     |   |          |   |    |    |                |      |    |                  | 55 | Mn     | Manganese<br>25 |      | ဥ        | Technetium<br>43 | 186 | Be       | Rhenium<br>75     |     |    |                | 144                       | S             | Neodymium                |
|                                   |       |     |   |          |   |    |    |                |      |    |                  | 52 | ပ်     | Chromium<br>24  | 96   | Mo       | Molybdenum<br>42 | 184 | ≽        | Tungsten<br>74    |     |    |                | 141                       | Ā             | Praseodymium             |
|                                   |       |     |   |          |   |    |    |                |      |    |                  | 51 | >      | Vanadium<br>23  | 93   | Q<br>N   | Niobium<br>41    | 181 | <u>Б</u> | Tantalum<br>73    |     |    |                | 140                       | ဝီ            | Cerium                   |
|                                   |       |     |   |          |   |    |    |                |      |    |                  | 48 | F      | Titanium<br>22  | 91   | Ż        | Zirconium<br>40  | 178 | Ξ        | Hafhium<br>72     |     |    |                | _                         |               |                          |
|                                   |       |     |   |          |   |    |    |                |      |    |                  | 45 | Sc     | Scandium<br>21  | 89   | >        | Yttrium<br>39    | 139 | Га       | Lanthanum<br>57 * | 227 | Ac | Actinium<br>89 | מסייים ל                  | U SCIIGO      | selles                   |
|                                   |       | =   |   |          |   | თ  | Be | Beryllium<br>4 | 24   | Mg | Magnesium<br>12  | 40 | Ca     | Calcium<br>20   | 88   | Š        | Strontium<br>38  | 137 | Ва       | Barium<br>56      | 226 | Ra | Radium<br>88   | * F8_71 Lanthanoid corios | - A 0+in 0 id | T 90-103 Actinoid series |
|                                   |       | _   |   |          |   | 7  | =  | Lithium<br>3   | 23   | Na | Sodium<br>11     | 39 | ¥      | Potassium<br>19 | 85   | Вb       | Rubidium<br>37   | 133 | Cs       | Caesium<br>55     | 223 | ቷ  | Francium<br>87 | * 58_71                   | 1 - 1 - 00 +  | T 30-100                 |
| 2013                              | 3     |     |   |          |   |    |    |                |      |    |                  |    | 50     | 70/21           | /N/I | 1/1      | ว                |     |          |                   |     |    |                |                           |               |                          |

| -                          |               |                    |               |                 |                 |                 |              |                 |                   |                   |                | •                  |                 |                   |
|----------------------------|---------------|--------------------|---------------|-----------------|-----------------|-----------------|--------------|-----------------|-------------------|-------------------|----------------|--------------------|-----------------|-------------------|
| opripo pripo               | 140           | 141                | 144           | 147             | 150             | 152             | 157          | 159             | 162               |                   |                | 169                | 173             | 175               |
| and series                 | S             | ሗ                  | P             | Pm              | Sm              | Ш               | gg           | Тb              | ò                 | 운                 | ш              | H                  | ΛÞ              | Ľ                 |
|                            | Cerium        | Praseodymium       |               | Promethium      | Samarium        | Europium        | Gadolinium   | Terbium         | Dysprosium        | 5                 | Erbium         | Thulium            | Ytterbium       | Lutetium          |
| Г                          | 28            | 29                 | 00            | 10              | 29              | 63              | 64           | 65              | 99                | /9                |                | 69                 | /0              | /1                |
| a = relative atomic mass   | 232           | 231                | 238           | 237             | 244             | 243             | 247          | 247             | 251               | 252               |                | 258                | 259             | 260               |
| X = atomic symbol          | 丘             | Ра                 | <b>-</b>      | Ν               | Pu              | Am              | Cm           | 쓢               | ర                 | Es                |                | Md                 | 2               | ۲                 |
| b = atomic (proton) number | Thorium<br>90 | Protactinium<br>91 | Uranium<br>92 | Neptunium<br>93 | Plutonium<br>94 | Americium<br>95 | Curium<br>96 | Berkelium<br>97 | Californium<br>98 | Einsteinium<br>99 | Fermium<br>100 | Mendelevium<br>101 | Nobelium<br>102 | Lawrencium<br>103 |
|                            |               |                    |               |                 |                 |                 |              |                 |                   |                   |                |                    |                 |                   |

Key

The volume of one mole of any gas is 24dm3 at room temperature and pressure (r.t.p.).