


UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Ordinary Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		



CHEMISTRY 5070/22

Paper 2 Theory

October/November 2013
1 hour 30 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Section A

Answer all questions.

Write your answers in the spaces provided in the Question Paper.

Section B

Answer any three questions.

Write your answers in the spaces provided in the Question Paper.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

A copy of the Periodic Table is printed on page 16.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This document consists of 16 printed pages.



Section A

For Examiner's Use

Answer **all** the questions in this section in the spaces provided.

The total mark for this section is 45.

A1 Choose from the following elements to answer the questions below.

chlorine
hydrogen
iron
lithium
nickel
nitrogen
oxygen
potassium
silver
sulfur
vanadium

zinc

Each element can be used once, more than once or not at all.

Which element

(a)	is liberated at the anode when an aqueous solution of potassium sulfate is electrolysed,
	[1]
(b)	is used as a catalyst in the manufacture of margarine,
	[1]
(c)	is a non-metallic solid, an atom of which contains only six valency electrons,
	[1]
(d)	is higher than sodium in the reactivity series,
	[1]
(e)	is in Period 5 of the Periodic Table,
	[1]
(f)	forms a white oxide which is amphoteric?
	[1]

[Total: 6]

A2 Carboxylic acids are a homologous series containing the –CO₂H group. The table shows some properties of the first four carboxylic acids in the series.

For Examiner's Use

carboxylic acid	molecular formula	density in g/cm ³	boiling point in °C
methanoic acid	CH ₂ O ₂	1.220	101
	C ₂ H ₄ O ₂	1.049	118
propanoic acid	C ₃ H ₆ O ₂	0.993	141
butanoic acid	C ₄ H ₈ O ₂	0.958	165

(a)	(i)	Describe how the density of these carboxylic acids varies with the number of carbon atoms in the molecule.
		[1]
	(ii)	Name the carboxylic acid with the molecular formula ${\rm C_2H_4O_2}.$
		[1]
	(iii)	Draw the structure of propanoic acid, showing all atoms and bonds.
		[1]
(b)		next carboxylic acid in this homologous series is pentanoic acid. tanoic acid has five carbon atoms.
	(i)	Deduce the molecular formula for pentanoic acid.
		[1]
	(ii)	Suggest a value for the boiling point of pentanoic acid.
		°C [1]
(c)	Buta	anoic acid, $C_3H_7CO_2H$, reacts with sodium to form a salt and a gas.
	(i)	Name the gas.
		[1]
	(ii)	Give the formula of the salt.
		[1]

For Examiner's Use


[Total: 11]

(d)		ers are formed when carboxylic acids react with alcohols. reaction is catalysed by hydrogen ions.
	(i)	Describe and explain the effect of a catalyst on reaction rate.
		[0]
		[2]
	(ii)	State one commercial use of esters.
		[1]
	(iii)	The structure of an ester is shown below.
		O H H H
		Name this ester.

А3	Silic	con is an element in Group IV of the Periodic Table.					
	(a)	Give the electronic configuration for a silicon atom.					
						[1]	
	(b)	Silic	con has three naturally o	ccurring isotopes.			
		Cor	nplete the following table	for two of these isotop	es.		
			isotope	²⁸ Si	³⁰ Si		
			number of protons				
			number of electrons				
			number of neutrons			[3]	
	(c)	Silic	con reacts with chlorine c	on heating to form silico	$\operatorname{en}(\operatorname{IV})$ chloride, $\operatorname{SiC} l_{\scriptscriptstyle A}$.	[0]	
	()		nstruct an equation for thi	•	4		
			·				
						[1]	
	(d)	Silicon(IV) chloride is a simple molecular compound.					
		(i)	Suggest two physical p	roperties of silicon(IV)	chloride other than solu	bility.	
			1				
			2			[2]	
		(ii)	Draw a 'dot-and-cross' or You only need to show to				

For Examiner's Use (e) Silicon(IV) chloride reacts with water to form silicon(IV) oxide. Part of the structure of silicon(IV) oxide is shown below.

For Examiner's Use



Explain, in terms of structure a point.	and bonding, why silicon(IV	/) oxide has a very high	າ melting
			[2]

[Total: 11]

A 4	The	carb	oon cycle regulates the amount of carbon dioxide in the atmosphere.
	(a)	(i)	State two processes which release carbon dioxide into the atmosphere.
			1
			2[2]
		(ii)	Name one process which removes carbon dioxide from the atmosphere.
			[1]
	(b)	Car	bon dioxide is a greenhouse gas.
		(i)	What is the meaning of the term <i>greenhouse gas</i> ?
			[1]
		(ii)	Name another greenhouse gas and give a natural source of this gas.
			name
			source[2]
	(c)	Car	bon dioxide dissolves in water to form a weakly acidic solution.
			$CO_2(g) + H_2O(I) \iff HCO_3^-(aq) + H^+(aq)$
		(i)	What is the meaning of the term weak acid?
			[1]
		(ii)	Describe how you could measure the pH of this solution other than by using a pH meter.
			[2]
	(d)	wate	lium hydrogencarbonate, $NaHCO_3$, decomposes on heating to form a carbonate, er and a gas which turns limewater milky. In a struct an equation for this reaction.

[2] [Total: 11] For Examiner's Use

A 5		student reacts magnesium ribbon with excess hydrochloric acid. ne follows the course of the reaction by measuring the volume of gas produced against ne.				
	(a)	Wri	te the equation for the reaction of magnesium with hydrochloric acid.			
			[1]			
	(b)	(i)	On the axes below draw a sketch graph to show how the volume of gas produced during the reaction varies with time and label this line 'A'. Label the axes with the appropriate units.			
			[2]			
		(ii)	The student then carries out the experiment at a lower temperature. All the other conditions remain the same.			
			On the axes above draw another line to show how the volume of gas produced varies with time and label this line 'B'. [1]			
	(c)	Ma	gnesium reacts with carbon to form the compound magnesium carbide.			
		Cal	culate the percentage by mass of magnesium in magnesium carbide, ${\rm MgC_2}$.			
			rol			
			[2]			
			[Total: 6]			

Section B

For Examiner's Use

Answer three questions from this section in the spaces provided.

The total mark for this section is 30.

В6	Alu	miniu	um is extracted from purified bauxite by electrolysis.
	(a)		scribe how this electrolysis is carried out and construct equations for the reactions urring at both the anode and cathode.
			[4]
	(b)	Wh	at properties of aluminium make it useful for
		(i)	making aircraft,
		(ii)	making electricity cables[2]
	(c)	Alu	minium is high in the reactivity series.
		(i)	Explain why aluminium does not react with aqueous copper(II) sulfate.
			[2]
		(ii)	When a few drops of aqueous sodium chloride are added to a mixture of aluminium and aqueous copper($\rm II$) sulfate, a vigorous reaction occurs.
			$copper(II)$ sulfate + aluminium \rightarrow aluminium sulfate + $copper$
			What type of reaction is this?
			[1]
		(iii)	State the formula of aluminium sulfate.
			[1]
			[Total: 10]

B7 Ethene is an unsaturated hydrocark	'non	n
--	------	---

For
Examiner's
Hea

(a) What is the meaning of each of these terms?

unsaturated	
hydrocarbon	[2]

- (b) Ethene can be manufactured by cracking.
 - (i) State the conditions used for cracking.

(ii) Construct an equation for the cracking of tetradecane, $\rm C_{14}H_{30}$, to form ethene and one other hydrocarbon.

- (c) Alkenes such as ethene can undergo addition polymerisation.
 - (i) State one use of poly(ethene).

(ii) The diagram below shows a section of a polymer chain.

Deduce the structure of the monomer which is used to make this polymer.

[1]

(d) Ethanol can be manufactured by the catalytic addition of steam to ethene.

For Examiner's Use

$$C_2H_4 + H_2O \rightarrow C_2H_5OH$$

If the reactants are not recycled, only 5% of the ethene is converted to ethanol.

Calculate the mass of ethanol formed from 0.4 tonnes of ethene when only 5% of the ethene is converted to ethanol.

[1 tonne is 1 000 000 grams]

mass of ethanoltonnes [3]

[Total: 10]

For Examiner's Use

В8		en hy ablish		HI, is heated in a closed t	tube, the following dynamic o	equilibrium is
			2HI(g)	$\rightleftharpoons H_2(g) + I_2(g)$	$\Delta H = + 9.6 \text{ kJ/mol}$	
	(a)	Wh	at is meant by th	e term <i>dynamic equilibriui</i>	m?	
						[2]
	(b)		e table shows the 25°C and 450°C.	e concentrations of HI(g),	$ m H_2(g)$ and $ m I_2(g)$ in the equilib	
			substance	concentration at 25 °C / mol/dm ³	concentration at 450 °C / mol/dm ³	
			HI(g)	0.94	0.79	
			H ₂ (g)	0.033	0.11	
			$I_2(g)$	0.033	0.11	
		(i)	The tube has a	volume of 50 cm ³ .		
			Calculate the m	nass of hydrogen iodide in	the equilibrium mixture at 25	°C.
						[2]
		(ii)	Describe and eat 25°C and 45		e concentrations of reactant	and products
						[2]

		13	
(c)	On your diagram lal the products,	by profile diagram for the decomposition of hydrogen iodide. So the reaction, ΔH .	
	energy	reactants	
		progress of reaction	[2]
(d)	An aqueous solution	n of hydrogen iodide contains iodide ions.	

For
Examiner's
Use

Describe a test for iodide ior	ns.	
		[2]

[Total: 10]

В9	The	com	npounds ammonium nitrate and ammonium sulfate are both fertilisers.	For
	(a)	Exp	plain why farmers add these fertilisers to soils.	Examiner's Use
			[1]	
	(b)	Am	monium sulfate can be prepared by adding sulfuric acid to aqueous ammonia.	
		Cor	nstruct the equation for this reaction.	
			[1]	
	(c)	Exc	cess acidity in soils can be treated by adding calcium hydroxide.	
		(i)	Give the formula of the ion present in calcium hydroxide which causes it to be alkaline.	
			[1]	
		(ii)	Explain why adding calcium hydroxide causes loss of nitrogen from fertilisers such as ammonium nitrate, which have been previously added to the soil.	
			[2]	
	(d)	A st	tudent titrated 10.0 cm ³ of aqueous calcium hydroxide with hydrochloric acid.	
			$Ca(OH)_2(aq) + 2HCl(aq) \rightarrow CaCl_2(aq) + 2H_2O(l)$	
			equired 4.00 cm ³ of 0.0100 mol/dm ³ hydrochloric acid to neutralise 10.0 cm ³ of leous calcium hydroxide.	
		Cal	culate the concentration of the calcium hydroxide.	
			mol/ dm ³ [3]	

For Examiner Use	Describe how to obtain pure dry crystals of calcium chloride from an aqueous solution of calcium chloride.	(e)
	[2]	
	[Total: 10]	

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

DATA SHEET
The Periodic Table of the Elements

			a ⊾ F	A) C			. 5	_ 4\ =	م د د		10 = E
		0	4 He Helium	20 Ne Neon	40 Ar Argon	84	Krypton 36	131 Xe Xenon 54	222 Rn Radon 86		Lu Lutetium 71
		II		19 Fluorine	35.5 C1 Chlorine	80	Bromine 35	127 I lodine 53	210 At Astatine 85		Y b Ytterbium 70
				16 Oxygen	32 S Sulfur 16	62	Selenium 34	128 Te Tellurium	209 Po Polonium 84		169 Tm Thullum 69
		>		14 Nitrogen 7	31 P Phosphorus 15	75	AS Arsenic 33	122 Sb Antimony 51	209 Bi Bismuth 83		167 Er Erbium 68
		2		12 Carbon 6	28 Si Silicon		Ge Germanium 32		207 Pb Lead		165 Ho Holmium 67
		≡		11 B Boron 5	27 A1 Aluminium 13		Ga Gallium 31	115 In Indium	204 T (Thallium 81		162 Dy Dysprosium 66
ts							Zinc 30	Cadmium 48	201 Hg Mercury 80		159 Tb Terbium 65
The Periodic Table of the Elements						64	Copper 29	108 Ag Silver 47	197 Au Gold 79		157 Gd Gadolinium 64
e of the	Group					29	Nickel 28	106 Pd Palladium 46	195 Pt Platinum 78		152 Eu Europium 63
odic Tabl	ğ			1		29	Cobalt 27	103 Rh Rhodium 45	192 Ir Iridium 77		Smarium 62
he Peric			1 Hydrogen			56	Fe Iron	Ruthenium 44	190 Os Osmium 76		Pm Promethium 61
_						55	Mn Manganese 25	Tc Technetium 43	186 Re Rhenium 75		Nd Neodymium 60
						52	Chromium	96 Mo Molybdenum 42	184 W Tungsten 74		Pr Praseodymium 59
						51	V Vanadium 23	Niobium 41	181 Ta Tantalum 73		140 Ce Cerium 58
						48	Titanium	91 Zr Zirconium 40	178 Hf Hafnium 72		
						45	Scandium 21	89 ×	139 La Lanthanum 57 *	227 AC Actinium 89	oid series I series
		=		9 Be Beryllium	24 Mg Magnesium	40	Calcium 20	Strontium	137 Ba Barium 56	226 Ra Radium	Lanthanc 3 Actinoic
		_		7 Li Lithium	23 Na Sodium	39	Potassium	85 Rb Rubidium 37	133 Cs Caesium 55	223 Fr Francium 87	* 58–71 Lanthanoid series † 90–103 Actinoid series
2013							E070/00	/O/N/13			

series	140 Ce	141 P	144 N	147 Pm	150 Sm	152 Eu	157 Gd	159 Tb	162 Dy	165 H	167 Er	169 Tm	173 Yb
5	Cerium 58	Praseodymium 59	Neodymium 60		Samarium 62	Europium 63	Gadolinium 64	Terbium 65	Dysprosium 66	Holmium 67	Erbium 68	Thulium 69	Ytterbium 70
lative atomic mass	232	231	238	237	244	243	247	247	251	252	257	258	259
omic symbol	ᄕ	Ра	>	Ν	Pn	Am	CB	路	₽	Es	Fm	Md	8
omic (proton) number	Thorium 90	Protactinium 91	Uranium 92	Neptunium 93	Plutonium 94	Americium 95	Curium 96	Berkelium 97	Californium 98	Einsteinium 99	Fermium 100	Mendelevium 101	Nobelium 102

Key

260 Lr Lawrendium 103

The volume of one mole of any gas is 24dm³ at room temperature and pressure (r.t.p.).