

Cambridge International Examinations

Cambridge Ordinary Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		



CHEMISTRY 5070/22

Paper 2 Theory

May/June 2016

1 hour 30 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Section A

Answer all questions.

Write your answers in the spaces provided in the Question Paper.

Section B

Answer any three questions.

Write your answers in the spaces provided in the Question Paper.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

A copy of the Periodic Table is printed on page 20.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This document consists of 19 printed pages and 1 blank page.




Section A

Answer **all** the questions in this section in the spaces provided.


The total mark for this section is 45.

A1 Choose from the following polymers to answer the questions.


polymer A




polymer B



polymer C



polymer D




polymer E


polymer F

© UCLES 2016


polymer G



polymer H



polymer I



Each polymer can be used once, more than once or not at all.

(a)	Which two polymers are polyesters?	
	and	[1]
(b)	Which polymer is used to make both clingfilm and plastic bags?	
		[1]
(c)	Give the letter of an addition polymer	
	Give the letter of a condensation polymer	[1]
(d)	Give the letter of a polymer that is a saturated hydrocarbon.	
		[1]
(e)	Which polymer could be part of a protein?	
		[1]

[Total: 5]

A2	Нус	Iroge	n fluoride, HF, has a simple molecular structure. It is soluble in water.		
	(a) Suggest one other physical property of hydrogen fluoride.				
	(b)	Hyd	rogen fluoride dissociates in water to form dilute hydrofluoric acid.		
		(i)	Write an equation to show the dissociation of hydrogen fluoride.		
		(ii)	Explain why an acidic solution is formed when hydrogen fluoride dissociates in water.		
			[1]		
	(c)	Dilu	te hydrofluoric acid reacts with aqueous calcium hydroxide.		
			$2HF(aq) + Ca(OH)_2(aq) \rightarrow CaF_2(aq) + 2H_2O(I)$		
			at is the minimum volume, in cm 3 , of 0.150 mol/dm 3 Ca(OH) $_2$ required to react completely a solution containing 0.200 g of HF?		
			volume of $Ca(OH)_2(aq) = \dots cm^3$ [3]		

(d)	Magnesium reacts with fluorine to make the ionic compound magnesium fluoride.				
	(i)	Predict two physical properties of magnesium fluoride.			
		1			
		2[2]			
	(ii)	Explain, in terms of electrons, how a magnesium atom reacts with a fluorine molecule, F_2 , to make a magnesium ion and two fluoride ions.			
		[2]			
		[Total: 10]			

Α3	Esters are used as	food flavourings an	nd solvents.
$\boldsymbol{\tau}$	Lotoro are acca ac	iood navodinigo di	ia doivoitto.

(2)	Draw the structure of	f othyl mothanoato	showing all of the atom	e and all of the hande
(a)	Draw the structure o	t etnvi metnanoate.	snowing all of the atom	s and all of the bonds.

b)	Ethy	thyl ethanoate evaporates at room temperature.					
	(i)	What is meant by the term evaporation?					
		[1]					
	(ii)	A sample of ethyl ethanoate in a beaker is moved into a colder room.					
		Explain, in terms of the kinetic particle theory, why this results in a decrease in the rate of evaporation.					

[1]

(iii) The table shows some information about different esters.

name	structure	relative molecular mass (M _r)
methyl ethanoate	CH ₃ CO ₂ CH ₃	74
ethyl ethanoate	CH ₃ CO ₂ C ₂ H ₅	88
propyl ethanoate	CH ₃ CO ₂ C ₃ H ₇	102
butyl ethanoate	CH ₃ CO ₂ C ₄ H ₉	116
pentyl ethanoate	CH ₃ CO ₂ C ₅ H ₁₁	130

Which ester has the lowest rate of evaporation at room temperature and pressure?
Explain your answer.
[2]
[Total: 6]

A 4	Sulfuric acid is manufactured by the contact process.			
	(a)	State the conditions used in the contact process.		
		temperature		
		pressure		
		catalyst	[2]	
	(b)	In the contact process, sulfur dioxide reacts with oxygen.		
		$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$		
		Describe and explain the effect of increasing the concentration of oxygen on the raction.	te of this	
			[2]	
	(c)	The catalyst used in the contact process increases the rate of the reaction. Describe one other advantage of using a catalyst in an industrial process.		
			[1]	
	(d)	Sulfuric acid is used to make the fertiliser potassium sulfate, K ₂ SO ₄ .		
		Calculate the percentage by mass of potassium in this fertiliser.		

[2]

[Total: 7]

A5 The statements give some of the chemical properties of cobalt and its compounds.

- Cobalt does not react with cold water.
- Cobalt fizzes slowly with dilute hydrochloric acid.
- Cobalt does not react with aqueous zinc nitrate.
- Cobalt reacts with aqueous silver nitrate.
- Cobalt(II) oxide reacts with magnesium to form cobalt.

(a)	Use the informa	tion to help arrange the following metals in order of reactivity.	
		cobalt, magnesium, silver, sodium and zinc	
	most reactive		
	least reactive		[2]
(b)	Construct the ed	quation for the reaction between cobalt(II) oxide, CoO, and magnesium.	
			[1]
(c)	Predict what hap	opens when cobalt(II) carbonate is heated strongly.	
			[1]
(d)	Cobalt has a me	elting point of 1495 °C.	
	-	s of structure and bonding, why a metal such as cobalt has a high messe a labelled diagram in your answer.	elting
			[2]
(e)	The symbol for o	one isotope of cobalt is $^{57}_{27}$ Co.	
	Another isotope	of cobalt has a nucleon number of 59.	

[Total: 7]

[1]

Write its symbol.

.....

			10
A 6	Rive	er wa	ter contains dissolved minerals and gases.
	(a)	Car	bon dioxide is one of the gases dissolved in river water.
			w the 'dot-and-cross' diagram to show the bonding in a molecule of carbon dioxide. Only w the outer-shell electrons.
			[1]
	(b)		er water often contains dissolved compounds such as ammonium nitrate and calcium sphate.
		(i)	State one source of both of these compounds.
			[1]
		(ii)	Describe and explain the environmental effect of the presence of these dissolved compounds in river water.

(c)	River water is often purified for use as drinking water.				
	Des	scribe three processes involved in the purification of river water.			
	prod	cess 1			
	prod	cess 2			
	prod	cess 3			
			[3]		
(d)	Wat	ter has a low melting point and is neutral ($pH = 7$).			
	(i)	Explain why water has a low melting point.			
			[1]		
	(ii)	A pH meter can be used to confirm that water is neutral.			
		Describe another way in which a student can confirm that water is neutral.			
			[1]		
		[Tota	l: 10		

BLANK PAGE

Section B

Answer three questions from this section in the spaces provided.

The total mark for this section is 30.

The	formula of lead(II) nitrate is $Pb(NO_3)_2$.	
(a)	which is insoluble in water.	
		•••
		•••
		4
(b)		of
	Construct the ionic equation, with state symbols, for this reaction.	
		2
(c)	Aqueous lead(II) nitrate is electrolysed using graphite electrodes. Bubbles of colourless g are formed at both electrodes.	38
	(i) Identify the gas formed at each electrode.	
	negative electrode (cathode)	
	positive electrode (anode)	•••
		2
	(ii) Construct the equation for the reaction at the cathode.	
		1
(d)	On heating, lead(II) nitrate decomposes to form PbO, NO_2 and O_2 .	
	Construct the equation for this reaction.	
		1
	(a) (b) (c)	are formed at both electrodes. (i) Identify the gas formed at each electrode. negative electrode (cathode)

[Total: 10]

 $\textbf{B8} \quad \text{Cyclohexene, C}_{6}\textbf{H}_{10}, \text{ is a cycloalkene.}$

Cycloalkenes react in a similar way to alkenes.

cyclohexene

(a)	Cyc	elohexene is an unsaturated hydrocarbon.
	(i)	What is meant by the term <i>unsaturated</i> ?
		[1
	(ii)	What is meant by the term <i>hydrocarbon</i> ?
		[1
(b)	Cor	nstruct the equation for the complete combustion of cyclohexene.
		[1
(c)	Сус	elohexene reacts with bromine.
	This	s is an addition reaction.
	(i)	Write the molecular formula of the product of this reaction.
		[1
	(ii)	What would be observed in this reaction?
		[1

	15
(d)	Cyclohexene can be manufactured from hexane as shown in the equation.
	$C_6H_{14} \rightarrow C_6H_{10} + 2H_2$
	Calculate the mass of cyclohexene that can be made from 258 g of hexane. $[M_{\rm r} \ {\rm of \ cyclohexene} = 82]$
	mass of cyclohexene = g [2]
(e)	Another cycloalkene has the following percentage composition by mass.
	C, 88.2%; H, 11.8%
	(i) Use the percentage composition by mass to show that the empirical formula of this cycloalkene is ${\rm C_5H_8}.$

[2]

(ii) The cycloalkene has a relative molecular mass, $M_{\rm r}$, of 68.

Draw the structure of the cycloalkene, showing all of the atoms and all of the bonds.

[1]

[Total: 10]

RQ	Carbon	monoxide	reacts	with	hydrogen	in a	reversible	reaction
DJ	Carbon	IIIOIIOXIUE	reacis	willi	nyuruqen	ша	. I evel Sible	reaction.

$$CO(g) + 2H_2(g) \rightleftharpoons CH_3OH(g) \Delta H = -91 \text{ kJ/mol}$$

The reaction reaches an equilibrium if carried out in a closed container.

(a)	Explain, in terms of bond breaking and bond forming, why this reaction is exothermic.				
		[2]			
(b)	Whe	en one mole of methanol, CH ₃ OH, is formed, 91 kJ of energy is released.			
		culate the amount of energy released when 160 g of methanol is formed. of methanol = 32]			
		energy released =kJ [2]			
(c)	Pre	dict, with a reason, how the position of equilibrium of this reaction changes as the			
	(i)	pressure is increased at constant temperature,			
		[2]			
	(ii)	temperature is increased at constant pressure.			
	(ii)				

(d)	Methanol and compound X react together to form methyl butanoate.				
	(i)	Name X.			
		[1]		
	(ii)	The reaction is normally carried out using a catalyst.			
		Name a suitable catalyst for this reaction.			
		[1]		
		[Total: 10	ე]		

 ${\bf B10}~{\rm Manganese}({\rm IV})$ oxide, ${\rm MnO_2},$ can be used in the preparation of both chlorine and oxygen.

Reaction 1
$$MnO_2(s) + 4HCl(aq) \rightarrow Cl_2(g) + 2H_2O(l) + MnCl_2(aq)$$

$$\textbf{Reaction 2} \hspace{0.5cm} 2 \text{H}_2 \text{O}_2(\text{aq}) \hspace{0.1cm} \boldsymbol{\rightarrow} \hspace{0.1cm} \text{O}_2(\text{g}) \hspace{0.1cm} + \hspace{0.1cm} 2 \text{H}_2 \text{O}(\text{I})$$

In reaction 2 manganese(IV) oxide acts as a catalyst.

(a) Reaction 1 converts chloride ions into chlorine molecules.

Explain why this is an example of oxidation.	
	111

(b) Reaction 1 is investigated using different masses of MnO₂. The results are shown in the table.

volume of HC <i>l</i> /cm ³	concentration of HC <i>I</i> (aq) in mol/dm ³	mass of MnO ₂ used /g	volume of Cl ₂ formed at room temperature and pressure /dm ³		
100	1.0	1.74	0.48		
100	1.0	0.87	0.24		

Explain the difference in the volume of chlorine formed.
[2

(c) Reaction 2 is investigated using different masses of MnO₂. The results are shown in the table.

volume of H ₂ O ₂ (aq) /cm ³	concentration of H ₂ O ₂ in mol/dm ³	mass of MnO ₂ used /g	volume of O ₂ formed at room temperature and pressure /dm ³
100	1.0	1.74	1.20
100	1.0	0.87	

Predict the volume of oxygen, measured at room temperature and pressure, when 0.87g of MnO₂ is used. Write your answer in the table. [1]

(d) Chlorine is bubbled through aqueous iron(II) chloride to form iron(III) chloride.

	Explain, with the aid of equations, how aqueous sodium hydroxide can be used to distinguish between aqueous iron(II) chloride and aqueous iron(III) chloride.
	[
(e)	Describe the chemical test for chlorine.
	test
	observation
	[2

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will

[Total: 10]

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2016 5070/22/M/J/16

be pleased to make amends at the earliest possible opportunity.

The Periodic Table of Elements

Group	IIIA	2 He	helium 4	10	Se	neon 20	18	Ar	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	R	radon			
	IIΛ			6	ш	fluorine 19	17	Cl	chlorine 35.5	35	Ā	bromine 80	53	П	iodine 127	85	Αt	astatine -			
	IN			80	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	<u>L</u>	tellurium 128	84	Ъо	polonium	116		livermorium -
	>			7	z	nitrogen 14	15	₾	phosphorus 31	33	As	arsenic 75	51	Sp	antimony 122	83	B	bismuth 209			
	>			9	ပ	carbon 12	41	Si	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	В	lead 207	114	F1	flerovium
	≡			2	Ω	boron 11	13	Ρl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	<i>1</i> 1	thallium 204			
										30	Zu	zinc 65	48	පි	cadmium 112	80	Нg	mercury 201	112	ű	copemicium
										29	Cn	copper 64	47	Ag	silver 108	79	Au	gold 197	111	Rg	roentgenium -
										28	ī	nickel 59	46	Pd	palladium 106	78	പ	platinum 195	110	Ds	darmstadtium -
										27	රි	cobalt 59	45	뫈	rhodium 103	77	'n	iridium 192	109	Μ	meitnerium -
		- I	hydrogen 1							26	Fe	iron 56	44	R	ruthenium 101	9/	SO	osmium 190	108	Hs	hassium -
		Key								25	Mn	manganese 55	43	ည	technetium -	75	Re	rhenium 186	107	Bh	bohrium –
					loc	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
			atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	q	niobium 93	73	<u>Б</u>	tantalum 181	105	Op	dubnium —	
					ato	rela				22	ï	titanium 48	40	Zr	zirconium 91	72	Ξ	hafnium 178	104	잪	rutherfordium —
										21	Sc	scandium 45	39	>	yttrium 89	57–71	lanthanoids		89–103	actinoids	
	=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	Š	strontium 88	26	Ba	barium 137	88	Ra	radium -
	_			8	:=	lithium 7	1	Na	sodium 23	19	¥	potassium 39	37	8	rubidium 85	55	S	caesium 133	87	μ̈	francium -

71 Lu	lutetium 175	103	۲	lawrencium -
۶ ۲b				_
e9 Tm	thulium 169	101	Md	mendelevium —
68 Fr	erbium 167	100	Fm	fermium -
67 Ho	holmium 165	66	Es	einsteinium –
。 Dy	dysprosium 163	86	ర	californium -
65 Tb	terbium 159	62	Ř	berkelium -
² Gd	gadolinium 157	96	Cm	curium
63 Eu	europium 152	92	Am	americium
62 Sm	samarium 150	94	Pu	plutonium
61 Pm	promethium —	93	d	neptunium —
9 PN	neodymium 144	92	\supset	uranium 238
59 Pr	praseodymium 141	91	Ра	protactinium 231
Se Ce	cerium 140	06	Т	thorium 232
57 La	lanthanum 139	68	Ac	actinium -

lanthanoids

actinoids

The volume of one mole of any gas is $24\,\mathrm{dm}^3$ at room temperature and pressure (r.t.p.)