

CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge Ordinary Level

MARK SCHEME for the October/November 2014 series

5054 PHYSICS

5054/22

Paper 2 (Theory), maximum raw mark 75

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2014	5054	22

Section A

- 1 (a) limit of proportionality (not breaking point) B1
- (b) (i) 8.5 cm **cao** B1
- (ii) 7.1 – 7.3 cm B1
- (c) $4.0 \times (2.7/7.2)$ **or** $5.0 \times (2.7/9.0)$ **or** 1.5 (N) **or** read from graph **or** 11.2 (cm)
0.148 – 0.152 kg **or** 148 – 152 g C1
A1 [5]
- 2 (a) (i) Fd **or** 2.5×0.18 C1
0.45 Nm A1
- (ii) force not applied at right angles to the tap B1
- (b) long(er) distance needs small(er) force (for same moment) **or** inversely related/proportional B1 [4]
- 3 (a) $V_1 = p_2 V_2 / p_1$ **or** $p \propto 1/V$ B1
 $1.0 \times 10^5 \times (1.8/2.0) \times 10^7$ C1
 $9.0 \times 10^{-3} \text{ m}^3$ **or** 9000 cm^3 A1
- (b) (i) $(\rho =) m/V$ **or** $(0.30/9.0) \times 10^{-3}$ C1
 $33(.3333) \text{ kg/m}^3$ **or** $0.033(3333) \text{ g/cm}^3$ A1
- (ii) helium mass/weight small (fraction of total/mass of air included) **or** this includes the weight of the cylinder B1 [6]
- 4 (a) (i) heat gained from burning fuel/combustion **or** friction between moving parts/with air/road **or** from (radiation of) Sun B1
- (ii) heat lost to air/surroundings **or** by convection (currents) **or** exhaust/hot gases/fumes **or** from exhaust **or** heat emitted (by hot car) **or** by radiation B1
- (b) at start chemical energy decreases **or** at start chemical energy to gravitational/potential energy (of car) increases **or** at end of process kinetic energy (of car **or** air) increases B1
B1
B1 [5]

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2014	5054	22

5	(a)	downward curve of correct curvature from marked 90 °C	B1	
		horizontal line at marked 58 °C	B1	
		downward (asymptotic) curve of correct curvature to marked 23 °C	B1	
	(b)	H marked halfway (by eye) along an intermediate horizontal line	B1	
	(c)	(Q =) mL or 45×220 9900 J	C1 A1	[6]
6	(a)	(the molecules) move faster or have more kinetic energy or accelerate ignore vibrate faster	B1	
	(b) (i)	faster/energetic molecules escape average speed decreases or slower molecules remain	B1 B1	
		temperature depends on average KE or heat taken from runner OR liquid becomes gas/vapour latent heat needed or bonds broken heat taken from runner	B1	
	(ii)	water vapour blown away or surrounding air less humid	B1	[5]
7	(a) (i)	lasts longer or one cell can be replaced without switching off the circuit or less (internal) resistance or if one fails the others still work	B1	
	(ii)	1.5 V	B1	
	(b) (i)	($R =$) V/I or $1.5/0.075$ 20Ω or $1.5/(0.075 - 6.0)$ 14Ω	C1 C1 A1	
	(ii)	decreases resistance of wire increases	B1 B1	[7]
8	(a)	one label correct <u>and</u> not contradicted C, 1S and 1B all correct and clear <u>and</u> none contradicted	C1 A1	
	(b)	any three from: magnetic field (between poles) (coil/wire) cuts field/flux or field/flux cuts (coil/wire) or field/flux changes (electromagnetic) induction brushes rub against/in contact with rings	B3	
	(c)	(half) distance across screen or count divisions of/measure wavelength or the wavelength corresponds to one rotation half distance multiplied by time base setting	B1 B1	[7]

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2014	5054	22

Section B

- 9 (a) changing speed/velocity C1
change in speed/velocity/time constant **or** $(v-u)/t$ constant **or** constant/equal
rate of change of speed/velocity A1 [2]
- (b) (a vector quantity has) direction B1 [1]
- (c) (i) 1. X between $t \geq 0$ and $t < 10$ s B1
2. Y between $t > 20$ s and $t < 30$ s B1
3. Z between $t > 10$ s and $t < 20$ s **or** between $t > 30$ s and $t < 40$ s B1
- (ii) 1. **two** speed values from graph between 15 and 35 s (± 1 mm) C1
two corresponding time values from graph between 15 and 35 s
(± 1 mm) **or** $(a =)\Delta v/t$ C1
500 m/s² A1
2. $(F =) ma$ **or** 8.4×500 C1
4200 N A1
- (iii) 1. arrow labelled F perpendicular to surface of Earth B1
arrow labelled R opposite to direction of travel (by eye) from rock B1
2. speed changes **or** density/pressure of air changes **or** cross-sectional
area (of rock) changes B1
- (iv) it hits the ground/surface of the earth **or** stops **or** speed is zero B1 [12]
- [Total: 15]
- 10 (a) 3.0×10^8 m/s B1 [1]
- (b) (i) 1. decreases **cao** B1
2. no change **cao** B1
3. decreases **cao** B1
- (ii) 1. i correctly marked (to normal) B1
2. r correctly marked (to normal) B1 [5]
- (c) (i) $\sin i/\sin r = n$ **or** $\sin i/\sin r = 1.5$ C1
 $\sin 89/\sin r = 1.5$ **or** $\sin 89/1.5$ **or** 0.67(0.666565) C1
42° **or** 41.8025° A1
- (ii) i equal to/close to 90° $\sin i/\sin 45$ $\sin^{-1}(1/n)/\sin^{-1}(1/1.5)$
and r less than 45° = 1.5 **and** 41.8° B1
or **or**
 i never bigger than $\sin i > 1$ r not be more than c
89°/90° B1 [5]

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2014	5054	22

- (d) (i) (sin) $i = 0$ **or** ray enters directly/
 along normal/perpendicular **or** wavefront/light hits surface
 (sin) $r = 0$ **or** no refraction all together
 all slows down together
- (ii) correct reflection at bottom surface (by eye) M1
 second correct reflection at top and no refraction at either point A1 [4]

[Total: 15]

- 11 (a) same element **or** same number of protons/atomic number B1
 different/particular number of neutrons **or** nucleons B1 [2]

- (b) (i) 38 cao B1
 (ii) 52 cao B1 [2]

- (c) ${}_{39}^{90}\text{Y}$ **or** ${}_{39}^{90}\text{Y}$ and ${}^0_{-1}\beta$ B1
 ${}^0_{-1}\beta$ ${}_{39}^{90}\text{Y}$ and ${}_{-1}^0\beta$ B1 [2]

- (d) (i) 87/29 **or** 3 (half-lives) **or** $6.0 \times 10^8/8$ C1
 7.5×10^7 A1
 (ii) any detector B1
 corresponding detection method B1

detector	film	(solid-state) detector	GM-tube	ionisation chamber	scintillation counter	cloud chamber
detection	fogged	count/reading	count/reading	count/reading	count/reading	track seen

no reduction with paper **or** (use of) electric/magnetic field **or** describe pattern of track M1

complete reduction with aluminium/lead **or** correct deflection of track in electric/magnetic field **or** no other track A1

- (iii) 1. unpredictable **or** not be known in advance **or** no set time between emissions **or** fluctuates **or** not fixed **or** counts obtained varies B1
 2. any **two** from:
 direction/in space
 time
 which nucleus decays B2 [9]

[Total: 15]