

Cambridge O Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

186109808

PHYSICS 5054/21

Paper 2 Theory May/June 2022

1 hour 45 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Section A: answer all questions.
- Section B: answer two questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has 20 pages. Any blank pages are indicated.

DC (CE/JG) 305640/2 © UCLES 2022


[Turn over

Section A

Answer all the questions in this section. Answer in the spaces provided.

1 Fig. 1.1 shows a model of the human arm. The rubber band represents the muscle that moves part of the arm XY up.

A mass is suspended from XY, as shown in Fig. 1.2. The weight of section XY is negligible and the model is at rest.






Fig. 1.1

Fig. 1.2 (not to scale)

(a)	(i)	State two ways in which the dimensions of the rubber band change as the mass added to section XY.	
	(ii)	State the form of energy stored in the stretched rubber band.	[1]
(b)	(i)	State the principle of moments.	
			[2]

(ii)	Explain why the force that the rubber band exerts on section XY is larger than the weight of the mass.
	[1]
(iii)	The mass suspended from section XY in Fig. 1.2 has a weight of 4.0 N.
	Calculate the force that the rubber band exerts on section XY.
	force =[2]
(iv)	Explain how your answer to (b)(iii) is different if the weight of section XY is not negligible.
	[1]
	[Total: 9]

2 Fig. 2.1 shows a wind turbine.




Fig. 2.1

(a) The wind blows directly towards the turbine with a speed of $12\,\text{m/s}$.

In one second, 60 000 kg of air passes through the circular area swept out by the blades. Calculate the kinetic energy of this mass of air.

	Kinetic energy =[3
b)	A wind turbine releases no carbon dioxide into the atmosphere while generating electricity Compared with a coal-fired power station, this is an advantage of using a wind turbine.

(i)	State one reason why it is important to reduce the amount of carbon dioxide produced.
	[1]
(ii)	Suggest one advantage of using a coal-fired power station compared with a wind turbine.
	[1]

(c)		A coal-fired power station releases 0.96 kg of carbon dioxide when it generates 1.0 kW h of electrical energy.					
	(i)	Define the kilowatt-hour (kWh).					
		[1]					
	(ii)	Calculate the mass of carbon dioxide saved when the wind turbine has a power output of 2000 kW and operates for 12 hours.					
		mass =[1]					
(d)	Win	d energy is a form of renewable energy.					
	Stat	e the name of one other form of renewable energy.					
		[1]					
		[Total: 8]					

					6		
3	(a)	Def	ine latent heat	of fus	sion.		
							[2]
	(b)				100 °C is heated steadily until its temperature rea	ches 400°C.	
		The	melting point	of the	e metal is 250 °C.		
		(i)	On Fig. 3.1, s	ketch	a graph to show how the temperature of the meta	ıl changes with tin	ne.
				400 7]		
		ten	nperature/°C	350-			
				300-			
				250-			
				200-			
				150-			
				100-			
				50-			
				0	<u> </u> 0		
					•	time	[2]
					Fig. 3.1		[2]
		(ii)	A sample of a	a differ	rent metal Q has a greater latent heat of fusion th	an the sample of	P.
			P and Q are capacity.	metals	s with the same melting points and the samples	have the same he	eat
					repeated with the sample of Q. This sample is supper second as is supplied to the sample of P.	oplied with the sar	me
			Explain how t	the gra	aph of temperature against time for Q differs from	the graph in (b)(i	i).

[2]

[Total: 6]

4 Two parallel rays of light, one red and one blue, enter a glass prism.

Fig. 4.1 shows both rays of light before they enter the prism. The blue ray is also shown incident on a different side of the prism after passing through the prism.




Fig. 4.1 (not to scale)

The ray of blue light strikes the left side of the prism at an angle equal to its critical angle *c*.

- (a) (i) On Fig. 4.1, mark and label the angle of incidence i and the angle of refraction r for the blue light as it enters the prism.[1]
 - (ii) On Fig. 4.1, continue the path of the blue light after it strikes the left side of the prism. [1]
 - (iii) The refractive index of glass for red light is smaller than the refractive index for blue light.

On Fig. 4.1, draw the path of the red light as it travels in the prism and after it strikes the left side of the prism. [2]

(b)	(i)	State what is meant by the critical angle.
		[2

(ii) The refractive index of glass for blue light is 1.5.

Calculate the critical angle *c* for blue light in glass. Show your working.

[Total: 8]

5 Fig. 5.1 shows a simple d.c. motor used in a toy car.




Fig. 5.1

Terminals A and B are connected to a battery and the motor rotates.

(a) On Fig. 5.1, add labels inside the boxes to identify the parts of the motor indicated. [2]
(b) State what happens to the rotation of the coil when:

(i) the number of turns on the coil is increased
(ii) the magnetic field between the poles of the magnet is reversed.

(c) The power supply to the motor is switched on and off at a steady rate.

Fig. 5.2 shows how the speed of the toy car varies with time as a result of the power supply being switched on and off.




Fig. 5.2

(i)	Describe the motion of the toy car.
	[2]
(ii)	The voltage supplied to the motor is switched on for a longer period of time and off for a longer period of time, at a steady rate.
	Suggest what happens to the motion of the toy car.
	[1]
	[Total: 7]

6 (a) A light-dependent resistor (LDR) is used to sense the amount of light in a room.

Fig. 6.1 shows part of the circuit used. The LDR is not shown.




Fig. 6.1

- (i) Complete Fig. 6.1 by drawing the circuit symbol for an LDR between points P and Q. [1]
- Explain how the voltmeter reading changes as the amount of light falling on the LDR increases.

ro1
1/1

(b) Fig. 6.2 shows part of a circuit containing three resistors.




Fig. 6.2

The current in the 4.0Ω resistor is 2.4 A.

(i) Show that the current in the 5.0Ω resistor is 1.9 A.

[2] (ii) Calculate the potential difference (p.d.) between points X and Y.

p.d. =[2]

[Total: 7]

Section B begins over the page

Section B

Answer two questions from this section. Answer in the spaces provided.

7 Fig. 7.1 shows a toy helicopter. It can hover and travel through the air.




Fig. 7.1

A student flies the toy helicopter on a journey from A to B to C to D at a constant height.

Fig. 7.2 is a scale drawing of the path of the helicopter, viewed from above.




Fig. 7.2 (to scale)

Determine the total distance travelled by the toy helicopter.

(ii) The toy helicopter makes the journey in 40 s.Calculate its average speed during the journey.

© UCLES 2022 5054/21/M/J/22

(a) (i)

(b) (i)	State the difference between distance and displacement.	
(ii)	Determine the scale used to construct Fig. 7.2 and complete the sentence.	
	1 cm on the diagram represents on the path.	[1]
(iii)	Using Fig. 7.2, determine the size of the displacement of point D from point A.	
	Show your working.	
	size of displacement =	[2]
(iv)	Determine the angle between North and the direction of the displacement of point D from point A.	эm
	angle =	[1]
(v)	State what is meant by velocity.	
		[1]
(vi)	Another toy helicopter flies directly from point A to point D in 40 s.	
	Explain why the magnitude of the velocity of this toy helicopter is smaller than the answin (a)(ii).	ver
		[1]


)	it falls.
	Explain, in terms of the forces and acceleration, what happens as the helicopter falls and reaches terminal velocity.
	[5]

[Total: 15]

Question 8 begins over the page.

8 Fig. 8.1 shows a water manometer.

When the water manometer is connected to a sealed container of gas and the tap is opened, the water levels change, as shown in Fig. 8.2.



(a) Define pressure.

		[1]
(b)	(i)	Explain why the water levels are the same on both sides of the manometer in Fig. 8.1.
		[1]
	(ii)	Explain why the water levels change when the gas tap is opened with the manometer connected as in Fig. 8.2.
		[1]
	(iii)	Explain why the water levels stop changing at the levels shown in Fig. 8.2.

(c) The water in the manometer has a density of $1000\,\text{kg/m}^3$.

	Atm	ospheric pressure is $1.00 \times 10^5 \text{Pa}$ and the gravitational field strength g is 10m/s^2 .
	Calo	culate the pressure of the gas inside the sealed container.
	Give	e your answer to 3 significant figures.
		pressure =[4]
(d)	The	temperature of the gas inside the sealed container increases.
	Usir	ng ideas about molecules explain why the water levels change.
		[4]
(e)	This	s manometer is not suitable for measuring a gas pressure of $2.5 \times 10^5 \text{Pa}$.
	(i)	Explain why.
		[2]
	(ii)	State one change to this manometer that allows it to measure a gas pressure of $2.5 \times 10^5 \text{Pa}$ in a normal school laboratory.
		[1]
		[Total: 15]

9 Table 9.1 shows details of seven different nuclides.

Table 9.1

nuclide	radiation emitted	half-life
hydrogen-2	none	_
hydrogen-3	beta	12 year
francium-223	beta	22 min
iridium-192	gamma	74 day
phosphorus-32	beta	14 day
radon-222	alpha	4 day
technetium-99	gamma	6 hour

(a)	(i)	Hydrogen-2 and hydrogen-3 are isotopes of the element hydrogen.
		Define the term isotope.
		[2]
	(ii)	The equation for the decay of phosphorus-32 (P-32) as it emits a beta particle is:
		$^{32}_{15}P \rightarrow ^{0}_{-1}\beta + ^{32}_{16}Q$
		Explain whether the equation shows that Q is another isotope of phosphorus. Q is not the chemical symbol for the atom.
		[1]
(b)		e of the sources in Table 9.1 is used in a medical procedure to detect unusual bone ctures. It is injected into a patient and the radiation emitted is detected outside the body.
	(i)	State which source in Table 9.1 is most suitable for this type of medical use.
		[1]
	(ii)	Explain two reasons for your choice

(c)	don gas is one natural source of background radiation.		
	Son	ne causes of background radiation are man-made, for example, X-rays.	
	(i)	State one other natural source of background radiation.	
			[1]
	(ii)	State one other man-made source of background radiation.	[4]
	(iii)	State one harmful effect of background radiation.	[1]
(d)	One fusion reaction that occurs is:		
		^2_1H + ^3_1H \rightarrow ^4_2He + $^{}_{}X$	
	(i)	Complete the equation to show the missing proton and nucleon numbers.	[1]
	(ii)	Deduce the name of particle X.	
			[1]
	(iii)	Suggest where this fusion reaction takes place.	
			[1]
(e)	Cor	npare the properties of alpha-particles and beta-particles in terms of their:	
		 ability to penetrate through materials ionising effects deflection in a magnetic field. 	
	•••••		[3]
			[၁]

[Total: 15]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.