

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the May/June 2007 question paper

9701 CHEMISTRY

9701/05

Paper 5 (Planning, Analysis and Evaluation),
maximum raw mark 30

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2007 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme	Syllabus	Paper
	GCE A/AS LEVEL – May/June 2007	9701	05

Question	Sections	Indicative material	Mark	
1 (a)	PLAN Problem	(i) Uses collision theory to predict that the rate of formation of $H_2(g)$ increases as the concentration of HCl increases	[1]	
		(ii) Uses collision theory to explain how rate of reaction increases with increasing temperature	[1]	[2]
(b)	PLAN Problem	<u>Concentration</u> of HCl identified as independent variable <i>[HCl] is acceptable</i>	[1]	[1]
(c)	PLAN Problem	States that the (total) volume of solution must be kept constant, or States that the amount/size/length/mass/surface area of the magnesium ribbon must be kept constant	[1]	[1]
(d)(i)	PLAN Methods	Lists apparatus for the reaction of $Mg/acid$, collection <u>and measurement</u> of gas and timing gas collection <i>Connecting tube does not need to be listed gas could be measured by full test-tube etc. A diagram is acceptable if a timing device is mentioned in the text</i>	[1]	
(ii)		Dilutes a range of volumes of HCl sufficient for the experiment <i>A minimum of 5 different concentration solutions is required Total volume does not have to be constant</i>	[1]	
(iii)		Prepares diluted solutions using measuring cylinder, pipette or burette	[1]	
(iv)		Describes how collection of a stated volume of H_2 will be timed in each experiment, or Volume of H_2 collected in a stated time is described, or Volume of H_2 collected recorded at fixed intervals to enable graph to be plotted	[1]	
(v)		Reference to the way in which total volume being kept constant, or temperature kept constant, or way in which other variable from (c) is controlled	[1]	

Page 3	Mark Scheme	Syllabus	Paper
	GCE A/AS LEVEL – May/June 2007	9701	05

Question	Sections	Indicative material	Mark	
(vi)		Candidate selects a range of suitable volumes of acid or states a range of concentrations to be used <i>Volume of acid should cover the range from starting volume (concentration) to at least half the starting volume (concentration)</i> <i>Total volume must be constant <u>unless</u> a correct (relative) concentration has been given</i> <i>Ignore starting with a concentration of <u>less</u> than 2 mol dm⁻³ hydrochloric acid.</i>	[1]	
(vii)		<i>Do <u>not</u> accept concentrations greater than 2 mol dm⁻³</i> The plan is presented logically with an effective way of preventing loss of gas <i>The use of dropping funnels or thistle funnels is permitted for addition of acid without loss of gas</i>	[1]	[7]
(e)	PLAN Methods	Table has columns for volume of acid and volume of water, *** time (if fixed volume of gas is collected) or volume of gas (if gas collected after fixed time) <u>rate</u> *** <i>Candidates may tabulate concentration instead of volume of acid and volume of water BUT TO QUALIFY FOR THIS MARK they must have shown numbers (volume of acid and volume of water) when describing a dilution in the text</i> Each column shown has correct units Candidate explains the graph (<u>valid for the method described</u>) which is to be drawn or the calculation to be performed or how the volume of gas – collected at fixed time interval or time – for collection of a fixed volume of gas will provide information in support of or against the prediction in (a)(i) <i>Examiners will expect increased concentration/increased rate</i> or <i>larger volume in fixed time linked to higher concentration</i> <i>shorter time for fixed volume linked to higher concentration</i> <i>(or reverse argument)</i>	[1] [1] [1]	[3]
(f)	PLAN Methods	Candidate repeats the experiment keeping HCl constant and varying the temperature Description of how the temperature will be <u>controlled</u> is required	[1]	[1]

Page 6	Mark Scheme	Syllabus	Paper
	GCE A/AS LEVEL – May/June 2007	9701	05

Question	Sections	Indicative material	Mark	
(f)	ACE Conclusions	Supporting evidence must be given <u>from and fit the data plotted</u>	[1]	[2]
		Suitable experimental method: Refers to a straight line, (passing through the origin), with few points off the line or Experimental method not suitable: Reverse argument to above or Suitable experimental method: Experimental data gives a value of x that is very close to an integer or Experimental method not suitable: Experimental data does not give an integral value of x	[1]	
(g)	ACE Conclusions	Soluble silver salt named e.g. silver nitrate/ AgNO_3 <i>Accept $\text{Ag}^+(\text{aq})$, solution containing Ag^+ or solution containing silver(I)</i> <i>Do <u>not</u> accept Ag^+ or silver</i> or Soluble lead(II) salt named e.g. lead nitrate/ $\text{Pb}(\text{NO}_3)_2$ <i>Accept $\text{Pb}^{2+}(\text{aq})$, solution containing Pb^{2+} or solution containing lead(II)</i> <i>Do <u>not</u> accept Pb^{2+} or lead</i> If formula or cation is given it must be correct <i>Ignore any potential reaction of an anion in the reagent with Hg^{2+}</i>	[1]	[1]
			[Total: 15]	

Appendix

Data for Question 2

	A	B	C		D	E	F
expt	mass of beaker /g	mass of beaker + mercury chloride /g	mass of beaker + mercury /g		mass of mercury chloride /g	mass of mercury /g	mass of chlorine /g
					(B–A)	(C–A)	(B–C) (D–E)
1	54.87	55.52	55.30		0.65	0.43	0.22
2	54.64	55.88	55.59		1.24	0.95	0.29
3	56.70	58.38	57.94		1.68	1.24	0.44
4	51.03	53.34	52.53		2.31	1.50	0.81
5	55.33	58.74	57.84		3.41	2.51	0.90
6	53.05	57.20	56.10		4.15	3.05	1.10
7	53.92	58.57	57.17		4.65	3.25	1.40
8	55.26	61.09	59.57		5.83	4.31	1.52

Zero required as second decimal place. Treat each error as a separate error

Candidate plots the following masses:

y axis	x axis	equation
mercury	mercury chloride	slope x (201 + 35.5x) = 201
mercury chloride	mercury	slope x 201 = (201 + 35.5x)
chlorine	mercury chloride	slope x (201 + 35.5x) = 35.5x
mercury chloride	chlorine	slope x 35.5x = (201 + 35.5x)
mercury	chlorine	slope x 35.5x = 201
chlorine	mercury	slope x 201 = 35.5x