

**MARK SCHEME for the October/November 2011 question paper
for the guidance of teachers**

8780 PHYSICAL SCIENCE

8780/03

Paper 3, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.


Page 2	Mark Scheme: Teachers' version	Syllabus
	GCE AS LEVEL – October/November 2011	8780

- 1 (a) 8.0 – 9.5 (°C);
- (b) reversed scale
non-linear, high numbers closer, at least 4 and scale easy to use [1]
- [Total: 3]
- 2 (a) +3/3/III allow 3+ [1]
- (b) moles CO₂ produced = 15 [1]
 $V = nRT/p = \frac{(15 \times 8.31 \times 298)}{100 \times 10^3}$ correct conversion and substitution [1]
 0.37(1) m³ [1]
- [Total: 4]
- 3 (a) $W = 17\,200\text{ N}$, $F = 17\,200\text{ N}$ [1]
 (must use $g = 9.81$ or 9.8 Nkg^{-1})
- (b) (i) use of force/area $\rightarrow 17\,200/(2.4 \times 1.0)$ [1]
 7200 Pa (accept ecf) [1]
- (ii) use of $p = \rho g \Delta h$ [1]
 $\Delta h = 7200/(1080 \times 100) \rightarrow \Delta h = 0.67\text{ m}$ (accept ecf) [1]
- (c) mass of water displaced = $0.68 \times 1.0 \times 2.4 \times 1080 = 1760\text{ kg}$ [1]
- [Total: 6]


Page 3	Mark Scheme: Teachers' version	Syllabus
	GCE AS LEVEL – October/November 2011	8780

www.PapaCambridge.com

4 (a) (i) BF_3 drawn as trigonal planar



BF_4^- drawn as tetrahedral



[2]

allow [1] if two fully-correct dot-and-cross diagrams given in place of both structures

BF_3 named as trigonal planar

[1]

BF_4^- angle = $109\frac{1}{2}^\circ$

[1]

(ii) equal repulsion between 3 bonding pairs

[1]

(b) (i) dative/coordinate

[1]

(ii) lone pair donated from F^- to B

allow to BF_3

[1]

[Total: 7]

5 (a) (i) 1 mm – 1 m

[1]

(ii) recognition that it is a diffraction effect

[1]

radio waves wavelength much longer than microwaves / microwaves wavelength much less than size of mountain / radio waves wavelength similar to mountain

[1]

(b) (i) path difference for contributions from slits = n wavelengths
so waves in phase (and add)/constructive interference

[1]

[1]

(ii) path difference for contributions from slits = $[n + \frac{1}{2}]$ wavelengths

[1]

so waves out of phase (and subtract/cancel) / destructive interference

[1]

(iii) amplitude = maximum amplitude $\div \sqrt{2}$

[1]

(iv) 1. maxima and minima/fringes move further apart

[1]

2. maxima and minima/fringes move closer

[1]

[Total: 10]



Page 4	Mark Scheme: Teachers' version	Syllabus
	GCE AS LEVEL – October/November 2011	8780

- 6 (a) $\text{CH}_4 + \text{H}_2\text{O} \rightarrow \text{CO} + 3\text{H}_2$
- (b) (i) quotes/refers to data showing decreased yield as temp. increases
high temp. favours endothermic direction so forwards = exothermic [1]
- (ii) fewer molecules/moles on right, high pressure favours direction
producing fewer molecules (\therefore higher yield) [1]
- (iii) pressure is compromise between rate/yield and cost of maintaining high
pressure [1]
allow: pressure used is the maximum economic pressure / is the
highest economically viable pressure
- (c) (i) N_2 and H_2 have only (weak) induced dipole-induced dipole/van der
Waal forces of attraction, (strong) hydrogen bonding present between
 NH_3 molecules [1]
hydrogen bonding much stronger than induced dipole-induced dipole/
van der Waal forces (so more energy/higher temperature needed to
separate molecules) [1]
- (ii) cooling the mixture allows ammonia to be removed as a liquid [1]
allow a specific statement to the effect that ammonia is removed by
condensation
- (d) $\Delta H_f = [(-414.5) + 2(-81.0)] - [(-287.0) + (-320.5)]$ [1]
 $= 31 \text{ kJ mol}^{-1}$ [1]
- [Total: 10]**
- 7 (a) the hydrogen nucleus has less charge / smaller (not less mass) / lower speed [1]
- (b) (i) attempted use of momentum equation $\rightarrow 5 \times 0.4 = 3 \times 0.4 + 8m$ [1]
 $\rightarrow 2 \times 0.4 = 8m_B \rightarrow m = 0.10 \text{ kg}$ [1]
- (ii) KE before = $\frac{1}{2} \times 0.4 \times 5^2 = 5.0 \text{ J}$ OR KE after = $\frac{1}{2} \times 0.4 \times 5^2 + \frac{1}{2} \times 0.1 \times 8^2$ [1]
correct calculation for both (= 5 J) [1]
statement that kinetic energy before = kinetic energy after [1]
- [Total: 6]**

Page 5	Mark Scheme: Teachers' version	Syllabus
	GCE AS LEVEL – October/November 2011	8780

www.PapaCambridge.com


- 8 (a) (i) σ bonding involves end-on overlap of orbitals / clear diagram
 π bonding involves sideways overlap (of 'p' orbitals) / clear diagram
- (ii) diagram of ethene showing planar shape and π bond clearly drawn, e.g.



3 curly arrows correctly positioned [1]
 correct intermediate bromocarocation [1]
 1,2-dibromoethane [1]

- (ii) induced dipole on Br_2 , caused by high electron density on $\text{C}=\text{C}$ bond [1]

- (c) (i) correct structure for 2-bromopropane – displayed formula expected but allow below as minimum detail: [1]



- (ii) alcohol [1]
 (iii) H^+ and $\text{K}_2\text{Cr}_2\text{O}_7$ and heat [1]
 (iv) propanone [1]

[Total: 11]

Page 6	Mark Scheme: Teachers' version	Syllabus
	GCE AS LEVEL – October/November 2011	8780

- 9 (a) positive background dough
electrons embedded
- (b) mark (i) and (ii) as one entity
 α -particle fired at gold foil [1]

three points, including at least one observation and one linked conclusion, from:
foil very thin/leaf
most go straight through*
* leads to mostly empty space
(very) small percentage deflected through large angles**
** leads to very small/massive nucleus [max 3]

- (c) (i) two from:
electrons in allowed orbits (accept orbitals/shells)
orbits 'radiationless'
fixed numbers in each orbit [max 2]
- (ii) group numbers = number of outer shell electrons [1]
period = number of shells [1]

[Total: 10]

- 10 (a) (i) $2I_2 - 8I_3 - 0 \times I_1 = 0 \rightarrow I_3 = 4I_2$ [1]
- (ii) $I_2 = 1.6 \text{ A}, I_3 = 0.4 \text{ A}$ [1]
- (b) $(1 - I_1 - I_2 = 0 \rightarrow 1 - I_1 - 1.6 = 0 \rightarrow =) - 0.6 \text{ A}$ (or could be done at point G) [1]
- (c) use of Kirchhoff's 2nd law around suitable loop [1]
 $E = 13.2 \text{ V}$ [1]

[Total: 5]

Page 7	Mark Scheme: Teachers' version	Syllabus
	GCE AS LEVEL – October/November 2011	8780

11 (a) (i) simplest ratio of atoms of each element in a compound/molecule

(ii)

<u>Na</u>	<u>Cl</u>	<u>O</u>
$\frac{21.6}{23}$	$\frac{33.3}{35.5}$	$\frac{45.1}{16}$

[1]

0.939 0.938 2.82

= 1 : 1 : 3
= NaClO₃

[1]

(b) (i) moles HCl = $21.70 \times 0.263/1000 = 5.71 \times 10^{-3}$ (mol) [1]

moles Q₂CO₃ = $0.571/2 = 2.85 \times 10^{-3}$ (mol) [1]

$M_r(Q_2CO_3) = 0.394/2.85 \times 10^{-3} = 138$ [1]

(ii) $A_r(Q) = [138 - 60]/2$, mark is for 60 [1]

= 39 so Q= K/potassium [1]

[Total: 8]