### **CAMBRIDGE INTERNATIONAL EXAMINATIONS**

Cambridge International Advanced Subsidiary and Advanced Level

## MARK SCHEME for the May/June 2015 series

# 9700 BIOLOGY

9700/43

Paper 4 (A2 Structured Questions), maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

S CAMPRIDCE

www.PapaCambridge.com

Page 2	Mark Scheme Sy.	per
	Cambridge International AS/A Level – May/June 2015 970 970	2
Mark sche	me abbreviations:	Cally
;	separates marking points	Or.
1	alternative answers for the same point	8
R	reject	i.c.
Α	accept (for answers correctly cued by the question, or by extra guidance)	On
AW	alternative wording (where responses vary more than usual)	7
<u>underline</u>	actual word given must be used by candidate (grammatical variants accepted)	

## Mark scheme abbreviations:

max indicates the maximum number of marks that can be given

or reverse argument ora


marking point (with relevant number) mp

error carried forward ecf

**AVP** alternative valid point (examples given as guidance)

Page 3	Mark Scheme	Sy. oer
	Cambridge International AS/A Level – May/June 2015	970

1 (a)



(b) large surface area (to get) more, light/carbon dioxide; A gas exchange I oxygen

thinness

small(er)/short(er)/reduced,  $\underline{\text{diffusion}}$  distance for gases **OR** fast(er)  $\underline{\text{diffusion}}$  of gases ; **A** named gas, either CO<sub>2</sub> or O<sub>2</sub>

1 mark only if both points made but not related to features in italics

[2]

[1]

[2]

- (c) (i) have chloroplasts/varying thickness of (cell) walls/no plasmodesmata;
  - (ii) water potential/ $\Psi$ , of (guard) cell(s), increases/becomes less negative ; water leaves cell(s) ;
    - (by) osmosis/down a water potential gradient ; I diffuses

(guard cell) becomes, flaccid/less turgid/AW;

[max 3]

- 2 (a) has more than one polypeptide; A FSH has  $2/\alpha$  and  $\beta$ , polypeptides R has four has, prosthetic group/non-protein part/carbohydrate/sugar; [max 1]
  - **(b)** 1 produce/make, monoclonal antibodies specific to (u-h)FSH/anti(u-h)FSH monoclonal antibodies;
    - 2 ref. to column/framework, for, attachment/immobilisation; **R** test strip
    - 3 urine, added to/flows past/passed over, antibodies;
    - 4 (so) allowing, hormone/(h)FSH, to bind (to monoclonal antibodies);
    - 5 treatment needed to release, hormone/(h)FSH (from monoclonal antibodies); I filtering [max 3]

		2.
Page 4	Mark Scheme	Sy. per
	Cambridge International AS/A Level – May/June 2015	970
		~

- (c) 1 sugars need to be added/glycosylation; A bacteria cannot modify protein
  - 2 needs, Golgi body/rough endoplasmic reticulum; A bacteria lack, Golgi/rough endoplasmic reticulum
  - 3 ref. to problems in bacteria with, introns/wrong promoter/secretion/ora; [max
- (d) labels to correct recognisable structures

```
(secondary) oocyte; R ovum
zona pellucida;
corona radiata/cumulus oophorus;
fluid-(filled space)/antrum;
granulosa/follicle/follicular, cells;
```

theca; [max 3]

- (e) (i) comparison
  - 1 more mature follicles with r-hFSH; ora
  - 2 oestrogen (concentration), higher with r-hFSH; ora
  - 3 comparative data quote; e.g. 13 v 8 mature follicles

**OR** 6.55 v 3.95 nmol dm<sup>-3</sup> oestrogen concentration

**OR** manipulated figures

e.g. difference of 5/2.6 nmol dm<sup>-3</sup>/

62.5% increase (r) follicles / 65.8% (r) oestrogen

explanation

4 (because) r-hFSH, purer/more concentrated/ora

ÒR

(some) u-hFSH, damaged by extraction technique/degraded;

[max 4]

- (ii) 1 difference/difference described, is significant;
  - 2 not due to chance; **A** due to something other than chance
  - 3 smaller than, critical value/value for significance of, 0.05/5%; [max 2]

[Total: 14]

				20
Page 5		(	Mark Scheme S Cambridge International AS/A Level – May/June 2015	y. A. O. Der 970 O. Day
(a)	1	self	f-pollination <b>ora</b> for cross-pollination netes/alleles/genes/DNA, come(s) from one parent;	970 ADACAMBRIDA
	2	give	es, less genetic variation/more genetic uniformity;	36
	3	resi	ults in inbreeding;	•
	4	incr	reases homozygosity/decreases heterozygosity;	[max 3]
(b)	ant	hers	and stigma/stamens and carpels, closer together;	[1]
(c)	1	ran	ge of flower size in original population ;	
	2	gen	netic variation (affecting flower size) in original population ; I mutati	ion
	3	cha	inge in environment/selection pressure, is absence of, bees/insect pollination (in greenhouse);	
	4	plar	nts with small, flowers/petals, are, selected for/reproduce/at a selective advantage ; <b>ora</b>	
	5	<u>alle</u>	les for small size passed to offspring ; ora I gene	
	6	freq	quency of, advantageous/smallness, allele increases; ora	
	7	dire	ectional selection;	
	8	tem	perature/irrigation/space/competition, different in field and glass	house;
	9	sma	all size explanation linked to factor in mp8;	[max 5]
				[Total: 9]
(a)	(i)	1	habitat loss/urbanisation/roads/agriculture; R deforestation	
		2	human damage (to plants); e.g. trampling/camping/picking	
		3	climate change ; e.g. drought/storms	
		4	soil erosion;	
		5	loss of pollinators;	
		6	use of herbicides;	
		7	competition with/eaten by, introduced species;	
		8	pollution;	[max 2]

Page 5

3

4

Page 6	6		Mark Scheme Sy. 75	per
J			Mark Scheme Sylvandridge International AS/A Level – May/June 2015 970	2
	(ii)	1	to maintain biodiversity ;	Cambridge.
		2	to maintain, food chains/food webs/stability of ecosystems;	Tage
		3	to maintain, genetic diversity/genetic variation/gene pool;	
		4	resources (for humans); e.g. biofuel/food/medicines/wood	
		5	aesthetic reasons/(eco)tourism;	
		6	to maintain, nutrient cycle/soil structure/climate stability;	
		7	idea of ethical duty;	[max 3]
(b)	1	gib	oberellin moves (from embryo) to aleurone layer ;	
	2	ge	ne, switched on/transcribed/used to make mRNA;	
	3	am	nylase produced ; I released/stimulated	
	4	(ar	mylase), hydrolyses/digests, starch to maltose; I breaks down/converts/glud	cose
	5	for	, respiration/ATP/energy ;	
	6	for	, growth/development/cell division/mitosis, in embryo;	
	7	ΑV	P; e.g. role of, DELLA/PIF	[max 4]
(c)	(i)	1	survival: less risk of extinction (for high seed survival compared with low survival);	
		2	germination percentage: for low survival: as % germination increases, risk of extinction decreases;	
		3	for high survival: as % germination increases risk of extinction decreases until, 30–36 % germination, then risk of extinction increases;	
		4	use of paired figures; e.g. quote % germination and risk of extinction for ea high v low [mp1] 2 points on low survival line [mp2] 2 points on high survival line [mp3] allow ± one grid square for figures	ch of: [max 3]

D		,	Mark Sahama	
Pa	age 7		Mark Scheme Sy.  Cambridge International AS/A Level – May/June 2015 970	Do Oer
		(ii)	yes 1 (scraping) increases germination ;	Da Cambridge
			2 more germination lowers risk of extinction; <b>ora</b>	36
			3 if seeds don't survive long/for low survival value seeds, scraping is good	;
			<ul><li>no</li><li>4 if seeds do survive long-term/for high survival value seeds, a store of see remains in soil;</li></ul>	eds
			5 (avoid risk of) all germinating at once and perhaps all dying;	[max 3]
				[Total:15]
5	(a)	1	two (complete) sets of chromosomes/diploid/2n;	
		2	one of each chromosome, from each parent/maternal and paternal;	
		3	to allow (homologous) pairs to form during, meiosis/prophase 1/reduction div	rision; [max 2]
	(b)	mo	ost/high %/more than 70%, of females in <b>three</b> populations prefer calls from their own population ;	
		les	ss than half/44%, of females in, <b>one</b> population/population 60, prefer calls from their own population ; <b>ora</b>	[2]
	(c)	1	yes different chromosome numbers;	
		2	cannot interbreed to form fertile offspring/hybrids infertile;	
		3	(because) not all chromosomes will be able to pair in meiosis;	
		4	live in different, habitats/climatic regions  OR	
			geographical isolation;	
		5	(so) unlikely to interbreed/reproductively isolated;	
		6	most females prefer males from their own population; ora	
		7	differences in mating, call/behaviour;	
		8	no some females, willing to mate with/prefer, males from other populations;	
		9	phenotypically/morphologically, similar;	[max 4]
				[Total: 8]

[Total: 8]

		Cambridge International AS/A Level – May/June 2015 970 970	20
(a)	1	toxin may bind to receptors on postsynaptic (membrane);	Cambridge
	2	(so) stops ACh binding/inhibits depolarisation/ no action potentials/Na <sup>+</sup> ion channels stay shut ;	36
	3	(so) stimulates ACh receptors / causes (continuous) depolarisation / causes action potentials / opens Na <sup>+</sup> ion channels ;	
	4	reduces/stops, release/recycling, of ACh (by presynaptic neurone);	
	5	inhibits acetyl cholinesterase/AW; R denatures	[max 3]
(b)	1	enter, presynaptic neurone/AW;	
	2	causes vesicles (containing ACh);	
	3	to, move to/fuse with, (presynaptic) membrane;	
	4	(so) ACh released (into synaptic cleft)/exocytosis;	[max 3]
(c)	1	ensure one-way transmission;	
	2	filter out infrequent impulses/temporal summation; I weak	
	3	allow, interconnection/integration, of, nerve (cell) pathways/many neurones; <b>OR</b> spatial summation/convergence of impulses/divergence of impulses;	
	4	ref. memory/learning;	
	5	idea of inhibitory effect;	[max 2]

Mark Scheme

Page 8

6

[Total:8]

Р	age 9	Mark Scheme	Sy. per
		Cambridge International AS/A Level – May/June 2015	970
7	(a) 1	glucose phosphorylated by ATP;	Canno
	2	(forms) hexose/fructose, bisphosphate;	Tage
	3	raises energy level of/activates, glucose/sugar OR lowers activation energy of reaction;	COM

- 7 (a) 1 glucose phosphorylated by ATP;
  - 2 (forms) hexose/fructose, bisphosphate;
  - 3 raises energy level of/activates, glucose/sugar lowers activation energy of reaction;

breaks down to two TP;

- 5 6C  $\rightarrow$  2 × 3C;
- hydrogen (atoms) removed/dehydrogenated/oxidised;
- 2 reduced NAD formed; A NADH/NADH<sub>2</sub>
- ref. to 4 ATP produced/net gain of 2 ATP;
- pyruvate produced;
- 10 AVP; e.g. ref. to substrate level phosphorylation/dehydrogenase/ phosphofructokinase/hexokinase

[max 6]

(b)

	substrate level phosphorylation	oxidative phosphorylation	
enzymes are involved	✓	✓	
occurs in cytoplasm	✓	×	;
occurs in mitochondria	<b>√</b>	<b>√</b>	;
channel proteins are involved	×	<b>√</b>	;

	· · · · · · · · · · · · · · · · · · ·	32
Page 10	Mark Scheme	Syl Sylver
	Cambridge International AS/A Level – May/June 2015	970
(c)	seeds soaked in water	Carry
1	little/no, oxygen (in water) ;	Office
2	(mostly) anaerobic respiration;	Sylvan, Park per 970 Park Park Park Park Park Park Park Park
3	seeds after 12 hours in the soil (more) aerobic respiration / less anaerobic respiration;	•
4	mixture of substrates; e.g. 2 of carbohydrates, proteins and lipids	
5	seedlings after 21 days aerobic respiration ;	
6	substrate is, glucose/carbohydrate;	

7 ref. to presence of leaves/photosynthesis;

[Total:15]

[max 6]

www.PapaCambridge.com Page 11 **Mark Scheme** Cambridge International AS/A Level – May/June 2015

#### 8 (a) gene mutation;

a change in the, base(s)/nucleotide(s); e.g. base, substitution/deletion/addition

(b) parental genotypes

CC<sup>a</sup>Bb x C<sup>h</sup>C<sup>a</sup>Bb;

gametes

CB Cb CaB C<sup>a</sup>b C<sup>h</sup>B C<sup>h</sup>b C<sup>a</sup>B C<sup>a</sup>b; allow on Punnett square

offspring genotypes;; deduct one mark for each error max 1 ecf for offspring genotypes if only 4 given

offspring phenotypes;

phenotypes linked to genotypes;

	C <sup>h</sup> B	CªB	C <sup>h</sup> b	Cab
СВ	CC <sup>h</sup> BB	CC <sup>a</sup> BB	<b>CC<sup>h</sup>Bb</b> full black	CC <sup>a</sup> Bb
Cb	<b>CC<sup>h</sup>Bb</b> full black	CC <sup>a</sup> Bb full black	CC <sup>h</sup> bb full red	CC <sup>a</sup> bb full red
C <sup>a</sup> B	C <sup>a</sup> C <sup>h</sup> BB	CªCªBB albino black	C <sup>a</sup> C <sup>h</sup> Bb Him black	CªCªBb albino black
Cªb	C <sup>a</sup> C <sup>h</sup> Bb Him black	C <sup>a</sup> C <sup>a</sup> Bb albino black	<b>C</b> <sup>a</sup> <b>C</b> <sup>h</sup> <b>bb</b> Him red	C <sup>a</sup> C <sup>a</sup> bb albino red

[6]

[Total:8]

Pa	age 1	2	Mark Scheme Sy.	per
	J		Mark Scheme Sy.  Cambridge International AS/A Level – May/June 2015 970	2
9	(a)	1	obtain mRNA from $\beta$ cells (of islets of Langerhans of pancreas) ;	Cambridge
		2	reverse transcriptase;	196
		3	make (single-stranded) <u>c</u> DNA;	
		4	DNA polymerase used to make cDNA double stranded;	
		5	sticky ends created; A description	
		6	(obtain) plasmids;	
		7	cut with restriction, endonuclease/enzyme; A named e.g. EcoR1	
		8	ref. complementary sticky ends ;	
		9	cDNA/insulin gene, mixed with plasmid;	
		10	<u>DNA ligase</u> ;	
		11	seals nicks in sugar-phosphate backbone ; R anneals	[max 8]
	(b)	1	(recombinant) plasmids mixed with bacteria;	
		2	(some) bacteria, take up plasmids/transformed;	
		3	heat shock/calcium chloride solution/Ca 2+ ions/electroporation;	
		4	to identify bacteria containing plasmids grow on, agar/medium, containing antibiotic (A); <b>A</b> ampicillin	
		5	plasmid contains, antibiotic (A)/ampicillin, resistance gene(s);	
		6	bacteria with plasmid survive ; ora	
		7	to identify recombinant bacteria replica plate; A description e.g. sponge/velvet pad/absorbent paper	
		8	(onto) agar/medium, containing second antibiotic (B); A tetracycline	
		9	$(tet^R/B/2^{nd})$ resistance gene inactivated (by insertion of new, DNA/gene)/AW;	
		10	(ID) colonies from, 1 <sup>st</sup> /ampicillin, plate that do not grow on, 2 <sup>nd</sup> /tetracycline, plate	e ; [max 7]

[Total:15]

			Cambridge International AS/A Level – May/June 2015 970	OS I
10	(a)	1	batch/penicillin nutrients, decrease/run out ;	OBC AMBridge
		2	so, secondary metabolite/penicillin, made;	36
		3	fermenters can be used (after cleaning) for different process;	
		4	if problem occurs only one batch affected;	
		5	needs little, monitoring/attention (once set up);	
		6	continuous/mycoprotein (fungus) kept in, exponential/log, phase (of growth);	
		7	(so) high, biomass/yield/production rate;	
		8	little/no, downtime;	
		9	small, vessels/space, required;	
		10	cost-effective;	[max 8]
	(b)	1	mouse is injected with an antigen;	
		2	wait for immune response to occur;	
		3	clonal selection ; A description e.g. antigen binds to, specific/virgin, B cell	
		4	clonal expansion; A description e.g. mitosis/division/cloning of B cells	
		5	B-lymphocytes/plasma cells, are extracted;	
		6	from the mouse's spleen;	
		7	fused with, cancer/myeloma/tumour, cells;	
		8	hybridoma cells formed;	
		9	hybridoma cells producing antibodies are identified;	

10 cultured on a large scale (to secrete monoclonal antibodies);

Mark Scheme

Page 13

[Total:15]

[max 7]