

Cambridge International AS & A Level

FURTHER MA	ATHEMATICS		9231/3
CENTRE NUMBER		CANDIDATE NUMBER	
CANDIDATE NAME			

Paper 3 Further Mechanics

May/June 2021

1 hour 30 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use 10 m s⁻².

INFORMATION


- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.

DC (CE/SW) 199008/2 © UCLES 2021

[Turn over






A uniform lamina ABCD consists of two isosceles triangles ABD and BCD. The diagonals of ABCD meet at the point O. The length of AO is AO

Find the distance of the centre of mass of the lamina from DB .	[3]

One end of a light elastic string of natural length $0.8 \mathrm{m}$ and modulus of elasticity $30 \mathrm{m}$ fixed point O on a smooth plane. The plane is inclined at an angle α to the horizonta A particle P of mass $2 \mathrm{kg}$ is attached to the other end of the string. The string lies alor slope of the plane with the particle below the level of O . The particle is projected w directly down the plane from the position where OP is equal to the natural length of	I, where $\sin \alpha = \frac{3}{5}$ ing a line of greates ith speed $\sqrt{2} \text{m s}^{-1}$
Find the maximum extension of the string during the subsequent motion.	[5]

[Turn over




Particles A and B, of masses 3m and m respectively, are connected by a light inextensible string of length a that passes through a fixed smooth ring R. Particle B hangs in equilibrium vertically below the ring. Particle A moves in horizontal circles on a smooth horizontal surface with speed $\frac{2}{5}\sqrt{ga}$. The angle between AR and BR is θ (see diagram). The normal reaction between A and the surface is $\frac{12}{5}mg$.

Find $\cos \theta$.	[3]

9231/33/M/J/21

© UCLES 2021 9231/33/M/J/21 **[Turn over**



A particle of mass m is attached to one end of a light inextensible string of length a. The other end of the string is attached to a fixed point O. The particle is initially held with the string taut at the point A, where OA makes an angle θ with the downward vertical through O. The particle is then projected with speed u perpendicular to OA and begins to move upwards in part of a vertical circle. The string goes slack when the particle is at the point B where angle AOB is a right angle. The speed of the particle when it is at B is $\frac{1}{2}u$ (see diagram).

Find the tension in the string at A , giving your answer in terms of m and g .	[8]
	•••••
	•••••
	•••••
	•••••
	•••••

 	• • • • • • • • • • • • • • • • • • • •




a)	Find an expression for v in terms of t , while P is moving upwards.



The displacement of P from O is x m at time t s.

Find an expression f	for x in terms of t ,	while P is moving	upwarus.	[2]
				••••••
Find. correct to 3 sig	nificant figures, t	he greatest height a	bove <i>O</i> reached by <i>P</i> .	[2]
Find, correct to 3 sig	gnificant figures, t	he greatest height a	bove <i>O</i> reached by <i>P</i> .	[2]
Find, correct to 3 sig	gnificant figures, t	he greatest height a	bove O reached by P .	[2]
Find, correct to 3 sig	gnificant figures, t	he greatest height a	bove <i>O</i> reached by <i>P</i> .	[2]
Find, correct to 3 sig	gnificant figures, t	he greatest height a	bove <i>O</i> reached by <i>P</i> .	[2]
Find, correct to 3 sig	gnificant figures, t	he greatest height a	bove O reached by P.	[2]
Find, correct to 3 sig	gnificant figures, t	he greatest height a	bove <i>O</i> reached by <i>P</i> .	[2]
			bove O reached by P.	

© UCLES 2021 9231/33/M/J/21 **[Turn over**



Two uniform smooth spheres A and B of equal radii have masses m and km respectively. Sphere A is moving with speed u on a smooth horizontal surface when it collides with sphere B which is at rest. Immediately before the collision, A's direction of motion makes an angle θ with the line of centres (see diagram). The coefficient of restitution between the spheres is $\frac{1}{3}$.

Show that the speed of <i>B</i> after the co	3(1+1/1)	

Given that $\tan \theta = \frac{1}{3}$, find	the value of k .	[6



[Turn over

(a)	Use the equation of the trajectory given in the List of formulae (MF19), together with the con $y = 0$, to establish an expression for the range R in terms of u , θ and g .	dition [2]
		•••••
		•••••
		•••••
		•••••
		•••••
		•••••
(b)	Deduce an expression for the maximum height H , in terms of u , θ and g .	[2]
(b)	Deduce an expression for the maximum height H , in terms of u , θ and g .	[2]
b)	Deduce an expression for the maximum height H , in terms of u , θ and g .	[2]
b)	Deduce an expression for the maximum height H , in terms of u , θ and g .	[2]
b)	Deduce an expression for the maximum height H , in terms of u , θ and g .	[2]
b)	Deduce an expression for the maximum height H , in terms of u , θ and g .	[2
b)	Deduce an expression for the maximum height H , in terms of u , θ and g .	[2
(b)	Deduce an expression for the maximum height H , in terms of u , θ and g .	[2]
(b)	Deduce an expression for the maximum height H , in terms of u , θ and g .	[2]
b)	Deduce an expression for the maximum height H , in terms of u , θ and g .	[2



It is	given that $R = \frac{4H}{\sqrt{3}}$.
(c)	Show that $\theta = 60^{\circ}$. [1]
It is	given also that $u = \sqrt{40} \mathrm{ms^{-1}}$.
	Find, by differentiating the equation of the trajectory or otherwise, the set of values of x for which
(u)	the direction of motion makes an angle of less than 45° with the horizontal. [4]



Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.			
		•••	
		•••	
© UCLES 2021	9231/33/M/J/21		



BLANK PAGE



BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2021 9231/33/M/J/21

