

Cambridge International AS & A Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

6 9 2 8 9 4 0 7 3 5

FURTHER MATHEMATICS

9231/42

Paper 4 Further Probability & Statistics

October/November 2021

1 hour 30 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has 16 pages. Any blank pages are indicated.

BLANK PAGE

equal size in country B. The results are summarised as follows.

The number, x, of pine trees was counted in each of 40 randomly chosen regions of equal size in country A. The number, y, of pine trees was counted in each of 60 randomly chosen regions of the same

1

Find a 95% cor	nfidence	interva	1 for the	difference	e hetweer	n the me	an num	her of nine	e trees in region	ים פו
this size in cou	ntries A	and B .	1 101 1110	difference	c between	ii tiic iiic	an num	oci oi pine	tices in region	[7]
	••••••	•••••	••••••	•••••		•••••				
										•••••
••••••	•••••	•••••	•••••	•••••		•••••		•••••	•••••	•••••
				•••••	•••••	•••••				•••••
										•••••
		•••••	•••••							•••••
		•••••								•••••
		•••••								•••••
		,								

2 It is claimed that the heights of a particular age group of boys follow a normal distribution with mean 125 cm and standard deviation 12 cm. Observations for a randomly chosen group of 60 boys in this age group are summarised in the following table. The table also gives the expected frequencies, correct to 2 decimal places, based on the normal distribution with mean 125 cm and standard deviation 12 cm.

Height, x cm	x < 100	$100 \leqslant x < 110$	$110 \leqslant x < 120$	$120 \leqslant x < 130$	$130 \leqslant x < 140$	$x \ge 140$
Observed frequency	0	3	15	23	11	8
Expected frequency	1.12	5.22	13.97	19.38	13.97	6.34

	•••••		•••••		•••••	
						• • • • • • • • • • • • • • • • • • • •
	•••••					• • • • • • • • • • • • • • • • • • • •
•••••	•••••					• • • • • • • • • • • • • • • • • • • •
Carry out a goo	dness of fit tes	st, at the 5% s	significance lev	vel, to determine	ne whether th	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determin	ne whether th	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determin	ne whether th	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determin	ne whether th	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determin	ne whether th	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determin	ne whether th	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determin	ne whether th	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determin	ne whether th	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determin	ne whether the	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determin	ne whether th	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determine	ne whether the	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determine	ne whether the	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determine	ne whether the	ne cla
Carry out a goo supported by the	dness of fit tes	st, at the 5% s	significance lev	vel, to determine	ne whether the	ne cla

	 •••••
 •••••	

 $\mathbf{3}$ The continuous random variable X has probability density function f given by

$$f(x) = \begin{cases} a + \frac{1}{5}x & 0 \le x < 1, \\ 2a - \frac{1}{5}x & 1 \le x \le 2, \\ 0 & \text{otherwise,} \end{cases}$$

where a is a constant.

Find the value	e of <i>a</i> .				
		••••••	 	 	
	•••••	•••••	 	 	•••••
			 	 	•••••
Find $E(X^2)$.				 	
Find $E(X^2)$.				 	
Find E(X ²).				 	
Find E(X ²).					
Find E(X ²).					
Find E(X ²).					

(c)	Find the cumulative distribution function of X . [3]

Test A Test B	46 36	32 28	29 49	12 37	33 48	18 35	25 41	40 31
(a) Carr diffe	y out a Wild rence in the	coxon rank population	-sum test at median ma	the 5% signrks obtaine	gnificance lo	evel to dete tests.	ermine whe	ther ther
•••••	•••••	•••••	•••••		•••••			

The interviewer considers using the given information to carry out a paired sample t-test to determine whether there is a difference in the population means for the two tests.

•••••	•••••	•••••	••••••		••••••	•••••		••••••	
	•••••		••••••		••••••	•••••		••••••	• • • • • • • • • • • • • • • • • • • •
					•••••			•••••	
					••••••	•••••			
				•••••			•••••		
	•••••	•••••		•••••		•••••	•••••		
	•••••			•••••		••••••	•••••		
		••••••		•••••		•••••	•••••		

(a)	Find the value of k .	[1]
(b)	Find the probability generating function $G_X(t)$ of X .	[2]
		•••••
		•••••
	random variable Y has probability generating function $G_Y(t) = \frac{1}{4} + \frac{1}{2}t + \frac{1}{4}t^2$.	
	random variable Z is the sum of X and Y .	. C 7
(c)	Assuming that X and Y are independent, find the probability generating function $G_Z(t)$ polynomial in t .	[3]
		•••••
		•••••

	12
(d)	Given that $E(Z) = \frac{13}{3}$, use $G_Z(t)$ to find $Var(Z)$. [3]

6

2.1	1.8	0.9	3.0	2.4	2.6	1.8	2.2	1.9	2.5
	ntist also of their m				12 fish o	of this type	e from lak	E = B, but	he only l
			$\sum y =$	= 24.48	$\sum y^2 = 5$	53.75			
the mean	e 10% sig mass of f st to be va	ish of this	level whe type in la	other the nake <i>B</i> . Yo	nean mass u should s	of fish of tate any as	this type s	in lake A is that you	s greater need to
• • • • • • • • • • • • • • • • • • • •									•••••
•••••		•••••	•••••			•••••	•••••	•••••	•••••
•••••				•••••					• • • • • • • • • • • • • • • • • • • •
•••••								•••••	•••••

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.						

15

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.