Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

2213454579

FURTHER MATHEMATICS **9231/13**

Paper 1 Further Pure Mathematics 1 **May/June 2022**

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

DC (CE) 214265/1

BLANK PAGE

1 **(a)** Sketch the curve with equation $y = \frac{x+1}{x-1}$. [2]

(b) Sketch the curve with equation $y = \frac{|x|+1}{|x|-1}$ and find the set of values of x for which $\frac{|x|+1}{|x|-1} < -2$. [4]

....

....

....

....

2 The cubic equation $x^3 + 5x^2 + 10x - 2 = 0$ has roots α, β, γ.

(a) Find the value of $\alpha^2 + \beta^2 + \gamma^2$. [3]

(b) Show that the matrix $\begin{pmatrix} 1 & \alpha & \beta \\ \alpha & 1 & \gamma \\ \beta & \gamma & 1 \end{pmatrix}$ is singular. [4]

3 A curve C has equation $y = \frac{ax^2 + x - 1}{x - 1}$, where a is a positive constant.

(a) Find the equations of the asymptotes of C. [3]

(b) Show that there is no point on C for which $1 < y < 1 + 4a$. [4]

....

....

....

....

....

....

....

....

....

....

(c) Sketch C. You do not need to find the coordinates of the intersections with the axes. [3]

4 Let $u_r = e^{rx}(e^{2x} - 2e^{x} + 1)$.

(a) Using the method of differences, or otherwise, find $\sum_{r=1}^{n} u_r$ in terms of n and x. [3]

....

....

....

....

....

....

....

....

....

....

....

....

....

(b) Deduce the set of non-zero values of x for which the infinite series

$$u_1 + u_2 + u_3 + \dots$$

is convergent and give the sum to infinity when this exists. [3]

....

....

....

....

....

....

....

....

....

....

(c) Using a standard result from the list of formulae (MF19), find $\sum_{r=1}^{n} \ln u_r$ in terms of n and x. [3]

5 Let $\mathbf{A} = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$, where a is a positive constant.

(a) State the type of the geometrical transformation in the x–y plane represented by **A**. [1]

....

....

(b) Prove by mathematical induction that, for all positive integers n,

$$\mathbf{A}^n = \begin{pmatrix} 1 & na \\ 0 & 1 \end{pmatrix}.$$ [5]

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

Let $\mathbf{B} = \begin{pmatrix} b & b \\ a^{-1} & a^{-1} \end{pmatrix}$, where b is a positive constant.

(c) Find the equations of the invariant lines, through the origin, of the transformation in the x–y plane represented by $\mathbf{A}^n\mathbf{B}$. [6]

6 The curve C has Cartesian equation $x^2 + xy + y^2 = a$, where a is a positive constant.

(a) Show that the polar equation of C is $r^2 = \dfrac{2a}{2 + \sin 2\theta}$. [3]

....

....

....

....

....

....

....

....

....

....

....

....

(b) Sketch the part of C for $0 \leqslant \theta \leqslant \frac{1}{4}\pi$. [2]

The region R is enclosed by this part of C, the initial line and the half-line $\theta = \frac{1}{4}\pi$.

(c) It is given that $\sin 2\theta$ may be expressed as $\dfrac{2\tan\theta}{1+\tan^2\theta}$. Use this result to show that the area of R is

$$\frac{1}{2}a\int_0^{\frac{1}{4}\pi} \frac{1+\tan^2\theta}{1+\tan\theta+\tan^2\theta}\,d\theta$$

and use the substitution $t = \tan\theta$ to find the exact value of this area. [8]

7 The position vectors of the points A, B, C, D are

$$7\mathbf{i}+4\mathbf{j}-\mathbf{k}, \qquad 11\mathbf{i}+3\mathbf{j}, \qquad 2\mathbf{i}+6\mathbf{j}+3\mathbf{k}, \qquad 2\mathbf{i}+7\mathbf{j}+\lambda\mathbf{k}$$

respectively.

(a) Given that the shortest distance between the line AB and the line CD is 3, show that $\lambda^2-5\lambda+4=0$. [7]

Let Π_1 be the plane ABD when $\lambda = 1$.

Let Π_2 be the plane ABD when $\lambda = 4$.

(b) **(i)** Write down an equation of Π_1, giving your answer in the form $\mathbf{r} = \mathbf{a} + s\mathbf{b} + t\mathbf{c}$. [2]

....

....

....

....

(ii) Find an equation of Π_2, giving your answer in the form $ax + by + cz = d$. [4]

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

(c) Find the acute angle between Π_1 and Π_2. [5]

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

BLANK PAGE

BLANK PAGE

BLANK PAGE