CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge Pre-U Certificate

MARK SCHEME for the May/June 2015 series

9791 CHEMISTRY

9791/04

Paper 4 (Practical), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.



Page 2	Mark Scheme	Syllabus	Paper
	Cambridge Pre-U – May/June 2015	9791	04

1 (a)	I Records in a table the volume of FA 1 , the volume of water and the reaction time for all five experiments.	[1]
	II Volumes given in cm ³ , time in seconds	[1]
	III All volumes recorded to 0.05, all times recorded to nearest second	[1]
	IV Selects 3 additional volumes of FA 1 , not less than 5.00 cm ³ and not closer together than 2.00 cm ³	[1]
	V All additional experiments have total volume of FA 1 + water = 20.00 cm ³	[1]
	VI Award if candidate T_{20}/T_{10} lies within 15% of supervisor T_{20}/T_{10} .	[1]
	VII Award if candidate T_{20}/T_{10} lies within 10% of supervisor T_{20}/T_{10} .	[1]
	VIII Award if $1.8 \le \text{candidate} T_{10} / T_{20} \le 2.2$	[1]
(b) (i)	Amount of $S_2O_3^{2-} = 1.5 \times 10^{-3} \text{ mol}$	[1]
	Concentration of iodine = $\frac{1}{2}(1.5 \times 10^{-3})/0.075 = 0.010 \text{ mol dm}^{-3}$	[1]
(ii)	Initial rates and initial concentrations correctly calculated for all five experiments.	[1]
(c)	I Initial rate on the <i>y-axis</i> with units of mol dm ⁻³ s ⁻¹ and initial concentration on the <i>x-axis</i> with units of mol dm ⁻³	[1]
	II Scales chosen to use more than ½ of each axis	[1]
	III All five points plotted as fine cross or encircled dot to within ½ small square and within the correct square	[1]
	IV Best-fit straight line drawn	[1]
(d) (i)	First order as graph is a straight line through the origin.	[1]
(ii)	rate = $k [H_2O_2] [I^-] [H^+]$	[1]
(iii)	Correctly calculates the gradient from the graph.	[1]
(iv)	Correctly calculates the initial concentration of H ₂ O ₂ as 0.0267 mol dm ⁻³ .	[1]
	Correctly calculates k (gradient = k [H ₂ O ₂] [H ⁺] k = gradient/(0.0267 × 0.276))	[1]
	Units of dm ⁶ mol ⁻² s ⁻¹	[1]
(v)	HSO_4^- is a weak acid and its degree of dissociation depends on the concentration of H^+ present in solution.	[1]
(e)	Times would be shorter/rate is faster so less accurate.	[1]
		[Total: 23]

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge Pre-U – May/June 2015	9791	04

2	FA 5 is CuCO ₃ + MnC 1 ₂	
(a) (i)	Off-white ppt.	[1]
(ii)	Fizzing and positive test for CO ₂ .	[1]
	Blue solution formed.	[1]
	Blue ppt on adding NaOH .	[1]
(b)	The soluble salt contains Mn ²⁺ . The insoluble salt contains Cu ²⁺ and CO ₃ ²⁻ . All three correct for two marks, two correct for one mark.	[2]
(c)	AgNO ₃ /NH ₃	[1]
	Notes problem of ppt with NH ₃ because of the presence of Mn ²⁺ . ALLOW answer which points out that you would need to filter off the silver halide ppt before testing with ammonia.	[1]
		[Total: 8]
3	FA 6 is Na ₂ SO ₃	
(a) (i)	White ppt and soluble in excess.	[1]
(ii)	White ppt and insoluble in excess.	[1]
(iii)	Purple solution turns to pale yellow/brown AND gives a brown ppt.	[1]
	Solution turns colourless on adding acid/ppt dissolves.	[1]
(iv)	Acidified KMnO₄ paper turns colourless.	[1]
	Gas identified as SO ₂ .	[1]
(b)	SO ₃ ²⁻	[1]
(c)	Sulfite has changed to sulfate.	[1]
	Oxidation by H ₂ O ₂ .	[1]
		[Total: 9]